

# Methane emissions from enteric fermentation as well as nitrogen and volatile solids excretions of German calves – a national approach

Ulrich Dämmgen\*, Ulrich Meyer\*\*, Claus Rösemann\*, Hans-Dieter Haenel\*, and Nicholas J. Hutchings\*\*\*

## Abstract

Enteric methane emission rates and (renal and faecal) nitrogen and volatile solids excretion rates were calculated using a procedure that reflects the development of the calves' rumen as well as German national animal performance data and representative diet properties.

Standard calves have a birth weight of 41 kg animal<sup>-1</sup>, a final weight of 125 kg animal<sup>-1</sup> and a mean weight gain of 0.67 kg animal<sup>-1</sup> d<sup>-1</sup>.

The emission rate of 9.4 kg methane per place and year and the methane conversion rate of 41 kJ MJ<sup>-1</sup> exceed those derived from expert judgements currently applied (4.4 kg place<sup>-1</sup> a<sup>-1</sup> and 20 kJ MJ<sup>-1</sup>, respectively). These differences are caused by different weights and weight gains. The newly derived methane emissions and methane conversion rates fall below those obtained from the application of IPCC Tier 2 procedures.

Nitrogen excretion rates amount to about 19 kg place<sup>-1</sup> a<sup>-1</sup>. The share of renally excreted nitrogen is 0.64 kg kg<sup>-1</sup> rather than 0.60 kg kg<sup>-1</sup> provided in the UNECE guidebook.

It is common practice to quantify volatile solids release rates according to IPCC guidance documents. However, this procedure had been shown to be inadequate. The results obtained in this work (144 kg place<sup>-1</sup> a<sup>-1</sup>) fall below those assessed with the IPCC methodology (about 500 kg place<sup>-1</sup> a<sup>-1</sup>).

**Keywords:** *calves, enteric fermentation, nitrogen, volatile solids, excretion, model*

## Zusammenfassung

### Methan-Emissionen aus der Verdauung sowie Stickstoff- und "volatile solids" – Ausscheidungen von deutschen Kälbern – ein nationaler Ansatz

Die Emissionsraten von Methan aus der Verdauung, die Ausscheidungsrate von (renalem und fäkalem) Stickstoff und von „volatile solids“ von Kälbern werden unter Berücksichtigung der Entwicklung des Kälberpensens sowie deutscher nationaler Leistungsdaten und repräsentativer Futtereigenschaften berechnet.

Standard-Kälber haben ein Geburtsgewicht von 41 kg, ein Endgewicht von 125 kg und eine mittlere Gewichtszunahme von 0,67 kg pro Tier und Tag.

Die Emissionsraten von 9,4 kg Methan pro Tierplatz und Jahr und eine Umwandlungsrate von 41 kJ MJ<sup>-1</sup> übersteigen die bisher verwendeten Raten (4,4 kg Tierplatz<sup>-1</sup> a<sup>-1</sup>) bzw. Methan-Umwandlungsrate (bisher 20 kJ MJ<sup>-1</sup>). Die Unterschiede sind auf unterschiedliche Gewichte und Gewichtszunahmen zurückzuführen. Die jetzt berechneten Methan-Emissionen und -Umwandlungsrate sind kleiner als die, die sich aus der Anwendung von IPCC Tier-2-Verfahren ergeben.

Die berechneten Stickstoff-Ausscheidungen belaufen sich auf etwa 19 kg pro Platz und Jahr. Der Anteil an renal ausgeschiedenem N von 0,64 kg kg<sup>-1</sup> ist höher als der im UNECE-Guidebook angegebene Wert von 0,60 kg kg<sup>-1</sup>.

Es ist üblich, die Ausscheidung von „volatile solids“ nach den Regeln des IPCC-Regelwerks zu berechnen. Das Verfahren ist jedoch unangemessen. Die in dieser Arbeit aus dem Futter berechneten Mengen (144 kg pro Platz und Jahr) liegen unter den Mengen, die sich bei Anwendung der IPCC-Methode ergäben (etwa 500 kg pro Platz und Jahr).

**Schlüsselwörter:** *Kälber, Verdauung, Stickstoff, volatile solids, Modell*

\* Johann Heinrich von Thünen Institute, Institute of Climate-Smart Agriculture, Bundesallee 50, 38116 Braunschweig, Germany

\*\* Friedrich Loeffler Institute (FLI), Federal Research Institute for Animal Health, Institute of Animal Nutrition, Bundesallee 50, 38116 Braunschweig, Germany

\*\*\* Aarhus University, Dept. of Agroecology, PO Box 50, Research Centre Foulum, 8830 Tjele, Denmark

Contact: ulrich.daemmgen@daemmgen.de

## 1 Introduction

Methane ( $\text{CH}_4$ ) from enteric fermentation in cattle contributes significantly to greenhouse gas emissions. International conventions aim at a reduction of these emissions. A first step is the adequate recording and reporting of these emissions. In order to ensure compatibility and comparability in international programmes, the United Nations Convention on Climate Change (IPCC) recommend the use of methodologies set down in the IPCC Guidelines for National Greenhouse Gas Inventories (IPCC, 1996, 2000, 2006)<sup>1</sup>. For emissions from enteric fermentation, the recommended procedure relates the  $\text{CH}_4$  emission rate to the gross energy (*GE*) intake rate, using a methane conversion rate (*MCR*). A detailed procedure to derive *GE* intake rates and *MCR* default values are provided in the Guidelines. For Western Europe the use of an *MCR* of 60 and 65 kJ MJ<sup>-1</sup> (or 6.0 and 6.5 %) is recommended in the 1996 and 2006 versions of the Guidelines, respectively. The application of default *MCR* explicitly includes “dairy cows and their young”. Although it is evident that *MCR* is a function of feed composition, poorer feed leading to higher *MCR* (see IPCC, 2006, pg. 10.30, comments on calves and feedlot cattle). However, no procedure to derive *MCR* reflecting this relation is explicitly recommended. For calves in particular, the fact that there is a delay after birth before ruminating becomes established is not taken into account.

This paper aims primarily at a derivation of a mean national  $\text{CH}_4$  emission rate (emission factor) and an *MCR* for German standard calves up to a weight of 125 kg animal<sup>-1</sup>. As the methodology used for these calculations presupposes the knowledge of the amounts of feed and of the feed ingredients' properties, the calculations are extended to derive consistent data for nitrogen (N) and volatile solids (*VS*) excretion rates.

## 2 Background

In emission inventories, animal subcategories are formed by populations receiving the same type of feed. In German animal production, it is usual to feed male and female calves up to a weight of 125 kg the same feeds. The respective time span lasts for about 18 weeks (or 125 days). The weight at birth is here taken to be 41 kg animal<sup>-1</sup>, the resulting average weight gain to be 0.67 kg animal<sup>-1</sup> d<sup>-1</sup>. Typical mean requirements for metabolizable energy (*ME*) during this time span are 21 to 24 MJ animal<sup>-1</sup> d<sup>-1</sup>. This data is in line with Stamer et al. (2004), DLG (1999) and KTBL (2010, pg 554).

As detailed data on the variation in time and region of weights and weight gains cannot be obtained, the calf described above serves as a **standard calf** in this work. Hence, the diet composition used to characterize energy and feed intake is a mean composition and does not reflect individual feeding situations.

The procedure to derive German national data describing the release of methane from enteric fermentation as well as of renal and faecal nitrogen (N) and volatile solids (*VS*) will comprise the following steps:

- assessment of calves' energy and nutrient requirements
- establishing diet composition and feed intake rates to meet these requirements
- listing feed properties needed
- calculating the respective release rates of methane
- calculating the excretion rates of N and *VS*

## 3 Feed requirements and intake rates

### 3.1 Metabolizable energy and nutrient requirements

Feed intake rates are governed by animal weights and weight gains and the resulting energy requirements. Germany uses the *ME* (metabolizable energy) system for the assessment of energy requirements of calves and for energetic feed evaluation. In Germany, official recommendations were provided in GfE (1997) and GfE (1999) to assess feeds satisfying the *ME* and nutrient requirements.

In the following, feed intake rates and diet compositions appropriate for German agriculture are used to calculate weekly *ME* intake rates. These are then compared to recommended intake rates that cover the span of a calf's life.

In addition the assessment of annual *ME* requirements presupposes the knowledge of animal rounds, i.e. the number of animals produced per place and year.

### 3.2 Feed intake rates and diet compositions

Feed intake rates and diet composition vary with the age of the animal. Examples can be derived from textbooks. Table 94 in Weiß et al. (2005) (similar in Meyer, 2005, and Teepker, 2007), supplemented with information from Kirchgeßner et al. (2008) formed the basis for the presumed diet in this paper. Typical weekly diet compositions are listed in Table 1.<sup>2</sup>

According to Kirchgeßner et al. (2008), hay should be of the best available quality, “tender” and “first cut”. This standard textbook in animal nutrition also lists typical concentrate compositions in calf feed (Table 2). An arithmetic mean composition of the four examples given in Kirchgeßner et al. (2008) was calculated and used in the subsequent calculations.

<sup>1</sup> For cattle, the information provided in IPCC (2000) does not deviate from IPCC (1996).

<sup>2</sup> Table 1 does not cover milk replacer. However, as milk replacer is used to substitute milk, the (known) properties and amounts of the milk fed are used instead.

**Table 1**

Amounts of milk, concentrates and hay fed to calves used in the calculations

week	colostrum	cows' milk	concentrates	hay	silage
kg animal <sup>-1</sup> d <sup>-1</sup>					
1	4.8	5			
2		6	0.2	0.1	
3		6	0.3	0.1	
4		6	0.4	0.1	
5		6	0.5	0.1	
6		6	0.7	0.1	
7		5	0.8	0.3	
8		5	0.9	0.5	
9		4.5	1.0	0.5	
10		4	1.2	0.6	
11		3.5	1.3	0.6	
12		2	1.4	0.5	0.5
13			1.5	0.5	1
14			1.5	0.5	1.5
15			1.5		2.5
16			1.5		3.5
17			1.6		4.5
18			1.6		5.5
kg animal <sup>-1</sup> round <sup>-1</sup>					
total	approx. 35	approx. 400	approx. 125	approx. 30	approx. 130

**Table 2**

Example concentrate compositions (Kirchgeßner et al., 2008)

diet constituent	I	II	III	IV	mean	unit
linseed expeller	0.20	0.12	0.10	0.00	0.105	kg kg <sup>-1</sup>
oat	0.10	0.00	0.00	0.18	0.070	kg kg <sup>-1</sup>
barley	0.20	0.00	0.29	0.20	0.173	kg kg <sup>-1</sup>
wheat	0.25	0.30	0.23	0.00	0.195	kg kg <sup>-1</sup>
maize	0.00	0.30	0.00	0.20	0.125	kg kg <sup>-1</sup>
wheat bran	0.09	0.08	0.00	0.00	0.043	kg kg <sup>-1</sup>
sugar beet pulp	0.00	0.00	0.13	0.08	0.053	kg kg <sup>-1</sup>
soya bean extraction meal	0.13	0.17	0.10	0.17	0.143	kg kg <sup>-1</sup>
rape seed extraction meal	0.00	0.00	0.12	0.14	0.065	kg kg <sup>-1</sup>
mineral feed	0.03	0.03	0.03	0.03	0.03	kg kg <sup>-1</sup>

### 3.3 Energy and relevant nutrient contents of feed ingredients

CH<sub>4</sub> emissions for fully ruminating animals are modelled using an approach suggested by Kirchgeßner et al. (1995) (for selection criteria see Dämmgen et al., 2012). The feed properties needed to calculate these emission rates, i.e. dry matter, crude fibre, N-free extracts, crude protein and crude fat contents were obtained from Beyer et al. (2004). They are collated in Table 3.

**Table 3**

Feed properties

feed ingredient	contents related to DM						
	dry matter	gross energy	metabolizable energy	crude fibre	N-free extracts	crude protein	crude fat
	unit	$\eta_{DM}$ kg kg <sup>-1</sup>	$\eta_{GE}$ MJ kg <sup>-1</sup>	$\eta_{ME}$ MJ kg <sup>-1</sup>	$\eta_{FI}$ kg kg <sup>-1</sup>	$\eta_{NFE}$ kg kg <sup>-1</sup>	$\eta_{XP}$ kg kg <sup>-1</sup>
colostrum <sup>a</sup>	0.15	23.98	18.55				
milk, 4.2 % fat	0.133	24.59	19.33				
hay	0.850	18.03	10.02	0.230	0.435	0.180	0.035
grass silage	0.350	18.40	10.20	0.245	0.452	0.162	0.042
maize silage	0.270	18.50	11.00	0.228	0.582	0.080	0.028
linseed expeller	0.900	20.69	12.70	0.110	0.375	0.380	0.370
oat	0.870	19.14	11.29	0.120	0.665	0.130	0.050
barley	0.870	18.56	12.91	0.050	0.765	0.135	0.025
wheat	0.870	18.60	13.44	0.020	0.485	0.115	0.025
maize	0.870	18.88	13.86	0.027	0.802	0.117	0.037
wheat bran	0.880	19.14	10.76	0.100	0.625	0.175	0.045
sugar beet pulp	0.920	16.28	12.66	0.200	0.645	0.095	0.005
soya bean extraction meal	0.900	19.96	14.00	0.080	0.345	0.495	0.017
rape seed extraction meal	0.900	20.30	12.20	0.130	0.350	0.370	0.020
minerals	1.000	0	0	0	0	0	0

<sup>a</sup> As the colostrum data set appeared to be inconsistent (see e.g. Foley and Otterby, 1978; Iváncsics and Kovács, 1999; and Morill et al., 2012) the energy contents of milk were used instead.

### 3.4 Comparison of calculated and recommended ME intake rates

As shown in Table 4, the use of the data listed in Tables 1 to 3 yields weekly ME input data whose period means fit the recommendations in GfE (1997), Table 8, for intervals of four to five weeks each, within the uncertainties involved. The animal weights are obtained from the birth weight of 41 kg animal<sup>-1</sup> and the constant weight gain of 0.666 kg animal<sup>-1</sup> d<sup>-1</sup>.

**Table 4**  
Metabolizable energy – requirements and recommendations

week	animal weight kg animal <sup>-1</sup>	calculated ME input MJ animal <sup>-1</sup> d <sup>-1</sup>	calculated mean ME MJ animal <sup>-1</sup> d <sup>-1</sup>	ME recommended MJ animal <sup>-1</sup> d <sup>-1</sup> (GfE, 1997)
1	45.7	12.78		
2	50.3	18.42	17.85	19
3	55.0	19.54		
4	59.6	20.66		
5	64.3	21.77		
6	69.0	24.01		
7	73.6	24.28	24.82	24
8	78.3	27.10		
9	83.0	26.94		
10	87.6	28.75		
11	92.3	28.59		
12	96.9	26.66	26.85	27.5
13	101.6	24.31		
14	106.3	25.96		
15	110.9	24.99		
16	115.6	28.27	30.47	30.5
17	120.3	32.68		
18	124.9	35.96		

### 3.5 Animal rounds

German official statistics report the numbers of animal places occupied at the census day (animal places or animal populations in IPCC terminology). In order to assess the emissions or excretions per place, the number of animals produced per place and year (i.e. the number of animal rounds) has to be taken into account.

The duration of a round covers both the time for the calves to grow to their final weight and the time for service time (i.e. for cleaning, drying and disinfection the animal place). The latter amounts to about of seven d round<sup>-1</sup> <sup>3</sup>. With a time span of 18 weeks for raising a calf, the number of animal rounds per year is thus about 2.8.

<sup>3</sup> In practice, service times range between one day and one month. The detailed instructions for FLI's experimental farm schedule give one week. This value is taken in subsequent calculations.

## 4 Calculation of methane emission rates

Calves only gradually emerge as ruminants (Kirchgeßner et al., 2008, pg. 430 ff). A pre-ruminant phase of about three weeks can be distinguished from a transition phase, during which the rumen develops its size and function. After about nine weeks calves can be considered functional ruminants. At that time, their typical MCR is that of cows (see Lockyer, 1997; Kurihara et al., 2002; Schönhusen et al., 2003; also Dämmgen et al., 2012).

No experimental data could be identified that describe CH<sub>4</sub> emission rates from calves in the transition phase.

### 4.1 General approach


#### 4.1.1 Physiological background

CH<sub>4</sub> is formed during the degradation of carbohydrates in the rumen. However, while concentrates and hay will enter the rumen, colostrum and milk bypass the rumen via the oesophageal groove and flow directly into the abomasum. The size and function of the rumen develop during the first seven to nine weeks of the calf's life (e.g. Baldwin et al., 2003).

The microbial degradation of carbohydrates presupposes a microbial community in the rumen that is adapted to the concentrates and the roughage fed. This community develops together with the rumen growth (e.g. Beharka et al., 1998).

#### 4.1.2 Stepwise calculation – an overview

These facts are accounted for in the following approach to characterize CH<sub>4</sub> emissions from calves in four steps where Steps 1 and 2 describe well defined situations, Steps 3 and 4 close the gap (Figure 1).



**Figure 1**  
Succession of steps used in the subsequent calculation procedure

**Step 1: Milk fed calves.** CH<sub>4</sub> emission rates for weeks one to three are zero, as the rumen has not developed. Hence, MCR is zero.

**Step 2: Fully ruminating calves with weights above 100 kg animal<sup>-1</sup>.** Kirchgeßner et al. (1995) describe the CH<sub>4</sub> emissions

from cattle for animals with a weight above 100 kg animal<sup>-1</sup>. Their methodology is applied to calculate the emission rates for weeks 15 to 18.

**Step 3: Fully ruminating calves with weights less than 100 kg animal<sup>-1</sup>.** It is assumed that the efficiency of CH<sub>4</sub> formation during weeks 9 to 15 is not substantially different from that of weeks 15 to 18. Hence it is characterized by the same methane conversion mechanisms, and that the emission rates obtained with the Kirchgeßner et al. (1995) approach can be extrapolated accordingly.

**Step 4: Transition from non-ruminant to ruminant calves.** It is assumed here that the gradual transition from non-ruminant to ruminant calves can be described using properties described in Step 3 combined with an appropriate efficiency parameter.

## 4.2 Calculation of methane emission rates of fully ruminating calves above 100 kg animal<sup>-1</sup> live weight (Step 2)

### 4.2.1 The method used for German dairy cows

In contrast to the IPCC methodology (IPCC, 1996, 2006), the German approach for dairy cows calculates CH<sub>4</sub> emission rates from feed intake rates and feed properties according to Dämmgen et al. (2012) (using the method proposed in Kirchgeßner et al., 1995) rather than from GE intake rates. The application of this method to calves was desirable.

The CH<sub>4</sub> excretion of the functional ruminant may be described by the approach of Kirchgeßner et al. (1995) (Equation (1)).

$$E_{\text{CH}_4, \text{K}} = a \cdot M_{\text{XFi}} + b \cdot M_{\text{NFE}} + c \cdot M_{\text{XP}} + d \cdot M_{\text{XF}} + e \quad (1)$$

where

$E_{\text{CH}_4, \text{K}}$	methane emission rate according to Kirchgeßner et al. (1995) (in kg animal <sup>-1</sup> d <sup>-1</sup> )
$a$	coefficient ( $a = 0.079$ kg kg <sup>-1</sup> )
$M_{\text{XFi}}$	intake rate of crude fibre (in kg animal <sup>-1</sup> d <sup>-1</sup> )
$b$	coefficient ( $b = 0.010$ kg kg <sup>-1</sup> )
$M_{\text{NFE}}$	intake rate of N-free extracts (in kg animal <sup>-1</sup> d <sup>-1</sup> )
$c$	coefficient ( $c = 0.026$ kg kg <sup>-1</sup> )
$M_{\text{XP}}$	intake rate of crude protein (in kg animal <sup>-1</sup> d <sup>-1</sup> )
$d$	coefficient ( $d = -0.212$ kg kg <sup>-1</sup> )
$M_{\text{XF}}$	intake rate of ether extract (crude fat) (in kg animal <sup>-1</sup> d <sup>-1</sup> )
$e$	constant ( $e = 0.063$ kg animal <sup>-1</sup> d <sup>-1</sup> )

and

$$M_{\text{XFi}} = M_{\text{DM}} \cdot \sum_i \eta_{\text{XFi}, i} \cdot x_i \quad (2)$$

where

$M_{\text{XFi}}$	intake rate of crude fibre (in kg animal <sup>-1</sup> a <sup>-1</sup> )
$M_{\text{DM}}$	dry matter intake rate (in kg animal <sup>-1</sup> a <sup>-1</sup> )
$\eta_{\text{XFi}, i}$	crude fibre content of feed constituent $i$ (in kg kg <sup>-1</sup> )
$x_i$	share of diet constituent $i$ (in kg kg <sup>-1</sup> )
etc.	

This approach is valid for cattle whose weights exceed 100 kg animal<sup>-1</sup>. Hence the feed composition for weeks 15 to 18 (inclusively) is used to estimate a mean daily CH<sub>4</sub> emission rate. This amounts to 0.032 kg animal<sup>-1</sup> d<sup>-1</sup> CH<sub>4</sub>.

### 4.2.2 Deriving a national methane conversion rate for fully ruminating calves

Within the IPCC methodology, it is customary to relate CH<sub>4</sub> emission rates as an energy loss to the GE intake rate using the methane conversion ratio  $MCR$ <sup>4</sup> which is defined as follows:

$$MCR = \frac{E_{\text{CH}_4} \cdot \eta_{\text{CH}_4}}{GE} \cdot \varepsilon \quad (3)$$

where

$MCR$	methane conversion ratio (in kJ MJ <sup>-1</sup> )
$E_{\text{CH}_4}$	methane emission rate (in kg animal <sup>-1</sup> d <sup>-1</sup> )
$\eta_{\text{CH}_4}$	energy content of methane ( $\eta_{\text{CH}_4} = 55,65$ MJ kg <sup>-1</sup> )
$\varepsilon$	energy units conversion factor ( $\varepsilon = 1000$ kJ MJ <sup>-1</sup> )
$GE$	gross energy intake rate (in MJ animal <sup>-1</sup> d <sup>-1</sup> )

For weeks 15 to 18, the mean daily GE intake rate amounts to 48.3 MJ animal<sup>-1</sup> d<sup>-1</sup> GE. Application of Equation (1) and the data provided in Tables 1 to 3 results in a mean CH<sub>4</sub> emission rate of 0.047 kg animal<sup>-1</sup> d<sup>-1</sup>, and an  $MCR$  of 54 kJ MJ<sup>-1</sup>. The value is lower than the ones proposed in IPCC (1996) and IPCC (2006) which is due to the comparatively high shares of concentrates. This results in lower  $MCR$  (see Kurihara et al., 2002; Schönhusen et al., 2003; the reasoning of the  $MCR$  of feedlot cattle in IPCC, 2006, pg 10.30; also Dämmgen et al., 2012).

### 4.3 Calculation of methane emission rates of fully ruminating calves below 100 kg animal<sup>-1</sup> live weight (Step 3)

Kirchgeßner's approach is not valid for animals whose weights fall below 100 kg animal<sup>-1</sup>. It is obvious from Equation (1) that constant  $e$  will dominate emissions for low feed intake rates. In order to close the gap, we assume that physiological conditions between week 9 and week 15 are similar to those in subsequent weeks, and apply the  $MCR$  determined for weeks 15 to 18 to the preceding weeks nine to fifteen. Therefore, the  $MCR$  of 54 kJ MJ<sup>-1</sup> obtained for weeks 15 to 18 is then applied to weeks nine to fifteen. GE intake rates are derived from ME requirements and feed properties, as listed in Tables 3 and 4.

### 4.4 Methane emissions during the transition period (Step 4)

The transition from a pre-ruminant newborn calf to a ruminant calf weighing 100 kg animal<sup>-1</sup> is described using an interpolation procedure. This is done by adding an effectiveness parameter  $P$  to describe rumen function, with  $P = 0$  for a newborn calf and  $P = 1$  for a fully functional ruminant:

<sup>4</sup>The IPCC terminology calls this entity methane conversion rate  $Y_m$ .

$$E_{\text{CH}_4, \text{trans}, j} = P_j \cdot E_{\text{CH}_4, \text{rum}} \quad (4)$$

with

$$E_{\text{CH}_4, \text{rum}} = E_{\text{CH}_4, \text{K}} \quad (5)$$

where

$E_{\text{CH}_4, \text{trans}, j}$  methane emission rate (factor) for week  $j$  in the transition period (in  $\text{kg animal}^{-1} \text{d}^{-1}$ )

$P_j$  effectiveness parameter for week  $j$  ( $0 \leq P \leq 1$ , dimensionless)

$E_{\text{CH}_4, \text{rum}}$  methane emission rate (factor) for a functional ruminant (in  $\text{kg animal}^{-1} \text{d}^{-1}$ ) as obtained by the application of Equation (1)

$E_{\text{CH}_4, \text{K}}$  methane emission rate according to Kirchgeßner et al. (1995) (in  $\text{kg animal}^{-1} \text{d}^{-1}$ )

Experimental data values for  $P$  are not available. Hence, assumptions have to be made for the amounts of  $P$  as a function of time. These are listed in Table 5. Here, variant 1 is a simple linear interpolation, whereas variant 2 may represent a more realistic approach that takes the non-linear growth of the rumen into account (Kirchgeßner et al., 2008; Coverdale et al., 2003; Pennstate College of Agricultural Sciences, 2011).

**Table 5**

Tentative time series of effectiveness parameters  $P_j$

week $j$	values of $P$	
	variant 1	variant 2
4	0.00	0.00
5	0.20	0.10
6	0.40	0.20
7	0.60	0.30
8	0.80	0.70
9	1.00	1.00

For each day in week  $j$ , the equivalents  $EE_{\text{CH}_4}$  of  $\text{CH}_4$  emission rates are assessed using the relation

$$EE_{\text{CH}_4, j} = GE_j \cdot P_j \cdot \frac{MCR_{\text{rum}}}{\varepsilon} \quad (6)$$

where

$EE_{\text{CH}_4, j}$  daily energy equivalent of  $\text{CH}_4$  emission rate in week  $j$  (in  $\text{MJ animal}^{-1} \text{d}^{-1}$ )

$GE_j$  GE intake rate in week  $j$  (in  $\text{MJ animal}^{-1} \text{d}^{-1}$ )

$P_j$  effectiveness parameter for week  $j$

$MCR_{\text{rum}}$  methane conversion rate for ruminating calves ( $MCR_{\text{rum}} = 54 \text{ kJ MJ}^{-1}$ )

$\varepsilon$  energy units conversion factor ( $\varepsilon = 1000 \text{ kJ MJ}^{-1}$ )

**Table 6**

GE intake rates, effectiveness parameter variants, energy equivalents of  $\text{CH}_4$  rates, and  $\text{CH}_4$  emission rates

week	GE intake rate	$P$		$EE_{\text{CH}_4}$		$E_{\text{CH}_4}$	
	$\text{MJ animal}^{-1} \text{d}^{-1}$	variant 1	variant 2	variant 1 $\text{MJ animal}^{-1} \text{d}^{-1}$	variant 2 $\text{MJ animal}^{-1} \text{d}^{-1}$	variant 1 $\text{kg animal}^{-1} \text{d}^{-1}$	variant 2 $\text{kg animal}^{-1} \text{d}^{-1}$
1	16.10	0.0	0.0	0.000	0.000	0.0000	0.0000
2	24.14	0.0	0.0	0.000	0.000	0.0000	0.0000
3	25.79	0.0	0.0	0.000	0.000	0.0000	0.0000
4	27.43	0.0	0.0	0.000	0.000	0.0000	0.0000
5	29.07	0.2	0.1	0.313	0.156	0.0056	0.0028
6	32.36	0.4	0.2	0.696	0.348	0.0125	0.0063
7	33.84	0.6	0.3	1.092	0.546	0.0196	0.0098
8	38.55	0.8	0.7	1.658	1.451	0.0298	0.0261
9	38.58	1.0	1.0	2.075	2.075	0.0373	0.0373
10	41.79	1.0	1.0	2.247	2.247	0.0404	0.0404
11	41.82	1.0	1.0	2.249	2.249	0.0404	0.0404
12	39.96	1.0	1.0	2.149	2.149	0.0386	0.0386
13	38.02	1.0	1.0	2.045	2.045	0.0367	0.0367
14	40.88	1.0	1.0	2.198	2.198	0.0395	0.0395
15	38.94	1.0	1.0	2.094	2.094	0.0376	0.0376
16	44.66	1.0	1.0	2.401	2.401	0.0432	0.0432
17	52.02	1.0	1.0	2.797	2.797	0.0503	0.0503
18	57.74	1.0	1.0	3.105	3.105	0.0558	0.0558
<b>Total</b>	<b><math>\text{MJ animal}^{-1}</math></b>			<b><math>\text{MJ animal}^{-1}</math></b>	<b><math>\text{MJ animal}^{-1}</math></b>	<b><math>\text{kg animal}^{-1}</math></b>	<b><math>\text{kg animal}^{-1}</math></b>
	4632			190	181	3.41	3.25

The total emission per animal for weeks 4 to 9 is obtained as follows:

$$E_{\text{CH}_4} = \sum_{j=4}^9 \left( \sum_{i=1}^7 \frac{EE_{\text{CH}_4, j}}{\eta_{\text{CH}_4}} \right) \quad (7)$$

where

$E_{\text{CH}_4}$  methane emission per animal (in kg animal<sup>-1</sup>)

$EE_{\text{CH}_4, j}$  daily energy equivalent of CH<sub>4</sub> emission rate in week j (in MJ animal<sup>-1</sup> d<sup>-1</sup>)

$\eta_{\text{CH}_4}$  energy content of methane ( $\eta_{\text{CH}_4} = 55,65 \text{ MJ kg}^{-1}$ )

$$EF_{\text{CH}_4} = E_{\text{CH}_4} \cdot n_{\text{round}} \quad (8)$$

where

$EF_{\text{CH}_4}$  CH<sub>4</sub> emission factor (in kg place<sup>-1</sup> a<sup>-1</sup> CH<sub>4</sub>)

$E_{\text{CH}_4}$  methane emission per animal (in kg animal<sup>-1</sup> d<sup>-1</sup>)

$n_{\text{round}}$  number of animal rounds (in animal place<sup>-1</sup> a<sup>-1</sup>)

For  $E_{\text{CH}_4} = 3.41 \text{ kg place}^{-1} \text{ a}^{-1}$  and  $n_{\text{round}} = 2.77 \text{ animal place}^{-1} \text{ a}^{-1}$ , the resulting emission factor for German standard calves is  $9.43 \text{ kg place}^{-1} \text{ a}^{-1} \text{ CH}_4$ .

It is mandatory within the reporting to UNFCCC to indicate respective *MCR* in the Common Reporting Format. Application of Equation (3) yields an *MCR* for calves of  $41.0 \text{ kJ MJ}^{-1}$  or 4.1 %.

#### 4.5 Predicted methane emissions by calves over a production period

Weekly input data are collated in Table 6. *GE* intake rates were obtained from a combination of figures provided in Tables 1 to 4. Using the two effectiveness parameter variants listed in Table 5 allows for the prediction of CH<sub>4</sub> emissions.

Variation 2 leads to slightly lower emissions, as would any variant that depicts a delayed (however likely) rumen development. In order to avoid underestimation of the emission rates, variant 1 should be used for reporting.

#### 4.6 Resulting national annual emission rates and methane conversion rates

The combination of emission per animal with the number of animal rounds yields the emission per place per year, the so-called emission factor  $EF_{\text{CH}_4}$ :

### 5 Nitrogen and volatile solids release rates

N and *VS* release rates are depending on feed intake, diet composition and – for N – on animal weight gained during the life span considered. IPCC (1996, 2006) as well as EMEP (2009) provide calculation procedures.

#### 5.1 Relevant feed properties

Feed properties are derived as weighted means from the relevant feed ingredients. Properties of the ingredients are taken from Beyer et al. (2004) (Table 7). The relevant amounts are listed in Tables 1 and 2, the relevant properties in Tables 3 and 7. For simplification, the amounts of colostrum are incorporated in the amounts of milk fed.

Table 7

Relevant feed properties for the calculation of N and VS excretion rates

		contents related to DM			
feed ingredient	unit	N content $X_N$ kg kg <sup>-1</sup>	digestibility of N $X_{\text{DN}}$ kg kg <sup>-1</sup>	ash content $X_{\text{ash}}$ kg kg <sup>-1</sup>	digestibility of OM $X_{\text{DOM}}$ kg kg <sup>-1</sup>
milk, 4.2 % fat		0.0411	0.95	0.073	0.98
hay		0.0288	0.70	0.120	0.73
grass silage		0.0259	0.63	0.100	0.71
maize silage		0.0128	0.45	0.080	0.74
linseed expeller		0.0608	0.85	0.065	0.79
oat		0.0208	0.74	0.035	0.73
barley		0.0216	0.74	0.025	0.86
wheat		0.0224	0.75	0.025	0.89
maize		0.0171	0.73	0.017	0.90
wheat bran		0.0280	0.75	0.055	0.71
sugar beet pulp		0.0088	0.50	0.050	0.90
soya bean extraction meal		0.0792	0.90	0.065	0.89
rape seed extraction meal		0.0592	0.84	0.080	0.77
minerals		0.0000	0.00	1.000	0.00

## 5.2 Annual nitrogen release rates

N release rates are obtained from the N mass balance:

$$m_{\text{excr}} = m_{\text{feed}} - m_{\text{g}} \quad (9)$$

$$m_{\text{excr, TAN}} = m_{\text{feed}} \cdot (1 - X_{\text{N}}) - m_{\text{g}} \quad (10)$$

where

$m_{\text{excr}}$  annual nitrogen release rate (in kg place<sup>-1</sup> a<sup>-1</sup> N)

$m_{\text{feed}}$  annual nitrogen intake rate with feed  
(in kg place<sup>-1</sup> a<sup>-1</sup> N)

$m_{\text{g}}$  amount of nitrogen retained in the animal  
(in kg place<sup>-1</sup> a<sup>-1</sup> N)

$m_{\text{excr, TAN}}$  total ammoniacal nitrogen (TAN) excretion rate  
(in kg place<sup>-1</sup> a<sup>-1</sup> N)

$X_{\text{N}}$  digestibility of nitrogen (in kg kg<sup>-1</sup>)

The N intake rate is the product of feed intake rate and N content of feed constituents for all constituents  $i$  and all time spans (weeks):

$$m_{\text{feed}} = n_{\text{round}} \cdot \sum_k \left( \sum_i m_{\text{feed, i, k}} \cdot X_{\text{N, i}} \right) \quad (11)$$

where

$m_{\text{feed}}$  annual nitrogen intake rate with feed  
(in kg place<sup>-1</sup> a<sup>-1</sup> N)

$n_{\text{round}}$  number of animal rounds (in a<sup>-1</sup>) (see Chapter 3.5)

$k$  running index of feeding days within one round  
( $1 \leq k \leq 126$ ) (for 18 weeks)

$m_{\text{feed, i, k}}$  daily intake rate of feed constituent  $i$  (dry matter)  
on day  $k$  (in kg d<sup>-1</sup>)

$X_{\text{N, i}}$  N content of feed constituent  $i$  (in kg kg<sup>-1</sup>)

The amount of **N taken in** with feed can be obtained by combining the information provided in Tables 1, 2 and 7 as expressed in Equation (11). The overall N input is 9.4 kg animal<sup>-1</sup> N or 26.0 kg place<sup>-1</sup> a<sup>-1</sup> N.

The amount of **N retained** per round is calculated from the weight gain (84 kg animal<sup>-1</sup>) and the mean N content of the weight gained per animal. GfE (1997) provide a mean protein content of the empty body of 182.2 g kg<sup>-1</sup>. The data base covers years between 1980 and 1990 (approximately). For the weight gain used in this work, Robelin and Chilliard (1989) investigating in Holstein Friesians found 193 g kg<sup>-1</sup>, Flachowsky et al. (1996) report 188 g kg<sup>-1</sup>. Janssen (2006) measured a mean protein content of German Holstein calves of 178.5 g kg<sup>-1</sup>. The "official" protein content of 182.2 g kg<sup>-1</sup> provided in GfE (1997), Table 1, was used in this work. With a mean N content of animal protein of 0.157 kg kg<sup>-1</sup>, a mean N content of 0.0287 kg kg<sup>-1</sup> N and a weight gain of 84 kg animal<sup>-1</sup> lead to an amount of N retained of 2.4 kg animal<sup>-1</sup> N or 6.7 kg place<sup>-1</sup> a<sup>-1</sup> N.

Overall N excretion rates consider intake rates and the amounts of N retained only. Renal N excretion rates are a function of the digestible N entering the animal metabolism (see Table 7).

The following N excretion rates are obtained

total N excretion	7.0 kg animal <sup>-1</sup> N	19.3 kg place <sup>-1</sup> a <sup>-1</sup> N
of which faecal	2.5 kg animal <sup>-1</sup> N	7.0 kg place <sup>-1</sup> a <sup>-1</sup> N
of which renal	4.4 kg animal <sup>-1</sup> N	12.3 kg place <sup>-1</sup> a <sup>-1</sup> N

The share of renally excreted nitrogen is 0.64 kg kg<sup>-1</sup> rather than 0.60 kg kg<sup>-1</sup> provided in the UNECE guidebook for "other cattle (young cattle, beef cattle, and suckling cows)" (EMEP, 2009, Table 3-8).

## 5.3 VS release rate

$VS$  release rates can be obtained from organic matter input rates and ash contents of the feed according to Equation (12) (Dämmgen et al., 2011)

$$VS = n_{\text{round}} \cdot \sum_k \left( \sum_i m_{\text{feed, i, k}} \cdot (1 - X_{\text{ash, feed, i}}) \cdot (1 - X_{\text{DOM, i}}) \right) \quad (12)$$

where

$VS$   $VS$  release rate (in kg place<sup>-1</sup> a<sup>-1</sup>  $VS$ )

$n_{\text{round}}$  number of animal rounds (in a<sup>-1</sup>) (see Chapter 3.5)

$k$  running index of feeding days within one round  
( $1 \leq k \leq 126$ ) (for 18 weeks)

$m_{\text{feed, i, k}}$  intake rate of feed constituent  $i$  (dry matter) on  
day  $k$  (in kg place<sup>-1</sup> d<sup>-1</sup>)

$X_{\text{ash, feed, i}}$  ash content of feed constituent  $i$  (in kg kg<sup>-1</sup>)

$X_{\text{DOM, i}}$  apparent digestibility of organic matter of feed  
constituent  $i$  (in kg kg<sup>-1</sup>)

A  $VS$  release rate of 52.1 kg animal<sup>-1</sup> or 144.1 kg place<sup>-1</sup> a<sup>-1</sup> is obtained for standard calves.

## 6 Discussion and conclusions

The description of emission rates from calves' enteric fermentation differs in principle from that of the release rates of nitrogen with faeces and urine and  $VS$  with faeces. The latter are approachable via balance considerations whereas enteric fermentation needs more sophisticated modelling. N excretion rates can be quantified in accordance with the UNECE guidance document (Dämmgen and Hutchings, 2008; EMEP, 2009), the calculations leading to CH<sub>4</sub> emission rates from enteric fermentation and  $VS$  release rates differ considerably from those recommended in the respective IPCC guidelines (IPCC 1996, 2006).

In this work, the **CH<sub>4</sub> emission factors  $EF_{\text{CH}_4}$**  and the **methane conversion rates  $MCR$**  are obtained using a combination of the cattle methane emission model (Kirchgeßner et al., 1995, see Dämmgen et al., 2012) for live weights above 100 kg animal<sup>-1</sup> and an adapted IPCC approach (using a national  $MCR$  for calves) for the animals between birth and a weight of 100 kg animal<sup>-1</sup>. Bridging the gap between the pre-ruminant calf and the fully ruminating calf requires assumptions that are plausible but nevertheless remain hypothe-

tical. In contrast, the IPCC approach of relating  $\text{CH}_4$  emission rates to  $GE$  intake rates is not applicable in principle (Dämmgen et al., 2012). Consequently, we recommend not using the IPCC default  $MCR$  values for cattle to quantify  $\text{CH}_4$  emission rates whenever the animal category “dairy cows and their young” (as in IPCC, 2006, Table 10.12) is divided in subcategories (calves, heifers, dairy cows). However, it would be appropriate to calculate emissions from energy demands, diet composition and feed properties using different  $MCR$  for calves on forage and on milk (as may be derived from IPCC, 2006, Table 10 A.2).

The average animal weights and weight gains listed in IPCC (2006), Table 10A.2, are not applicable to the German situation (see Chapter 2). Hence we favour a simple interpolation procedure to obtain adequate  $MCR$  for calves based on generally accepted principles, as performed in Chapter 3.5.2. Here, the  $MCR$  calculated for variant 1 (linear interpolation) exceeds those obtained potentially reflecting rumen growth (variant 2). Until such time that variant 2 is confirmed using empirical measurements, it is recommended to use the conservative results obtained with variant 1.

At present, the German agricultural emission inventory uses the IPCC Tier 2 methodology (with national energy intake rates and a national  $MCR$ ) to derive  $\text{CH}_4$  emissions from enteric fermentation of calves. Calculations are based on low weights at birth, high weight gains, a final weight of 100 kg animal<sup>-1</sup> and the application of an expert judgement of  $MCR$  of 20 kJ MJ<sup>-1</sup> (suggested by G. Flachowsky, Institute of Animal Nutrition, former Federal Agricultural Research Center, Braunschweig). It resulted in an emission factor ( $EF_{\text{CH}_4}$ ) of 4.3 kg place<sup>-1</sup> a<sup>-1</sup>  $\text{CH}_4$ . The methodology developed in this work describes calves with different properties, i.e. higher weights at birth (41 kg animal<sup>-1</sup> rather than 36 kg animal<sup>-1</sup>), a higher final weight (125 kg animal<sup>-1</sup> rather than 100 kg animal<sup>-1</sup>) and a lower daily weight gain (670 g animal<sup>-1</sup> d<sup>-1</sup> rather than 1066 g animal<sup>-1</sup> d<sup>-1</sup>). As shown in Chapter 3, it appears adequate to use an  $EF_{\text{CH}_4}$  of 9.4 kg place<sup>-1</sup> a<sup>-1</sup>  $\text{CH}_4$  and an  $MCR$  of 41 kJ MJ<sup>-1</sup> to describe the enteric emissions from German standard calves.

With regard to **N release rates with faeces and urine**, the value of 19.3 kg place<sup>-1</sup> a<sup>-1</sup> N obtained here is similar to the value previously used in German emission reporting (20.9 kg place<sup>-1</sup> a<sup>-1</sup> N; Haenel et al., 2012, Chapter 4.4.6.1). It contradicts the default value of 14 kg place<sup>-1</sup> a<sup>-1</sup> N provided in KTBL (2006), pg 428, for 4 animal rounds and a final weight of 100 kg animal<sup>-1</sup> (see reasoning in Haenel et al., 2012). The TAN content of 0.64 kg kg<sup>-1</sup> exceeds that previously used (0.60 kg kg<sup>-1</sup>) based on assumptions for cattle in general (Webb, 2001). This is attributed to the high digestibility of crude protein in milk and the high proportion of concentrates.

The modelling of **V/S release rates** differs from the IPCC approach in principle: the IPCC methodology is inadequate in general, as it relates  $V/S$  release rates to  $GE$  intake rates (Dämmgen et al., 2011). This is wrong in particular with (partly) milk fed calves as the zero fibre content of milk is not taken into account. The application of the IPCC (1996) methodology (IPCC, 1996, Table B-1: 1.46 kg animal<sup>-1</sup> d<sup>-1</sup>) to a standard German calf (using the default ash contents and di-

gestibilities of feeds provided) would yield a  $V/S$  release rate of about 500 kg place<sup>-1</sup> a<sup>-1</sup>. The difference between the IPCC approach and the results described in this work (144 kg place<sup>-1</sup> a<sup>-1</sup>) is substantial. Previous, less sophisticated German inventories made use of a  $V/S$  release rate of 209 kg place<sup>-1</sup> a<sup>-1</sup> (with animal performance data as listed above).

The emission factors  $EF_{\text{CH}_4}$  and the  $MCR$  as well as the excretion rates of  $V/S$  with faeces are based on standard descriptions of German calf production and derived using more realistic procedures than those provided in the IPCC guidance documents. The N excretion rates are calculated using more detailed data sets than before. Overall with respect to emission reporting, the new calculation procedures are considered to reflect more faithfully than previously the production conditions for the rearing of calves in Germany, and the accuracy of the description of calves has been improved.

We recommend using the new findings for future emission reporting.

## References

- Baldwin RL, McLeod KR, Klotz JL (2003) Rumen development, intestinal growth and hepatic metabolism in the pre-ruminant and postweaning ruminant. *J Dairy Sci* 87:E55-E65
- Beharka AA, Nagaraja TG, Morrill JL, Kennedy GA, Klemm RD (1998) Effects of form of the diet on anatomical, microbial, and fermentative development in the rumen of neonatal calves. *J Dairy Sci* 81:1946-1955
- Beyer M, Chudy A, Hoffmann L, Jentsch W, Laube W, Nehring K, Schiemann R (2004) Rostocker Futterbewertungssystem: Kennzahlen des Futterwertes und Futterbedarfs auf der Basis von Nettoenergie. Dummerstorf: Forschungsinst Biol landwirtschaftl Nutztiere, 392 p
- Coverdale JA, Tyler HD, Quigley JD, Brumm JA (2003) Effect of various levels of forage and form of diet on rumen development and growth in calves. *J Dairy Sci* 87:2554-2562
- Dämmgen U, Amon B, Gyldenkerne S, Hutchings NJ, Kleine Klausning H, Haenel H-D, Rösemann C (2011) Reassessment of the calculation procedure for the volatile solids excretion rates of cattle and pigs in the Austrian, Danish and German agricultural emission inventories. *Landbauforsch* 61(2):105-116
- Dämmgen U, Hutchings NJ (2008) Emissions of gaseous nitrogen species from manure management: a new approach. *Environ Pollut* 154(3):488-497
- Dämmgen U, Rösemann C, Haenel H-D, Hutchings NJ (2012) Enteric methane emissions from German dairy cows. *Landbauforsch* 62(1/2):21-31
- DLG-Deutsche Landwirtschaftsgesellschaft (1999) Leistungs- und qualitätsgerechte Junggründeraufzucht. Frankfurt a M: DLG, 19 p, DLG-Information 3/1999
- EMEP (2009) EMEP/EEA air pollutant emission inventory guidebook – 2009 [online]. To be found at <<http://www.eea.europa.eu/publications/emep-eea-emission-inventory-guidebook-2009>> [quoted 25.01.2013]
- Flachowsky G, Bugdol M, Löhnert HJ, Richter GH (1996) Einfluß unterschiedlicher Fütterungsintensität auf Lebendmasseentwicklung und Körperzusammensetzung von Aufzuchtälkälbern. *J Anim Physiol Anim Nutr* 76:9-21
- Foley JA, Terby DE (1978) Availability, storage, treatment, composition and feeding value of surplus colostrum: a review. *J Dairy Sci* 61:1033-1060
- GfE-Gesellschaft für Ernährungsphysiologie (1997) Empfehlungen zur Energieversorgung von Aufzuchtälkälbern und Aufzuchtrindern. *Proc Soc Nutr Physiol* 6:201-215
- GfE-Gesellschaft für Ernährungsphysiologie (1999) Empfehlungen zur Proteinversorgung von Aufzuchtälkälbern und Aufzuchtrindern. *Proc Soc Nutr Physiol* 8:155-164
- Haenel H-D, Rösemann C, Dämmgen U, Poddey E, Freibauer A, Döhler H, Eulich-Menden B, Wulf S, Dieterle M, Osterburg B (2012) Calculations of gaseous and particulate emissions from German agriculture 1990–2010:

- report on methods and data (RMD) Submission 2012. Braunschweig : vTI, 394 p, Landbauforsch SH 356
- IPCC-Intergovernmental Panel on Climate Change (1996) Revised 1996 IPCC guidelines for national greenhouse gas inventories : vol 3: Reference manual [online]. To be found at <<http://www.ipcc-nggip.iges.or.jp/public/gl/inv6.html>> [quoted 25.01.2013]
- IPCC-Intergovernmental Panel on Climate Change (2000) Good practice guidance and uncertainty measurement in national greenhouse gas inventories : chapter 4: Agriculture [online] To be found at <[http://www.ipcc-nggip.iges.or.jp/public/gp/english/4\\_Agriculture.pdf](http://www.ipcc-nggip.iges.or.jp/public/gp/english/4_Agriculture.pdf)> [quoted 25.01.2013]
- IPCC-Intergovernmental Panel on Climate Change (2006) 2006 IPCC guidelines for national greenhouse gas inventories : vol 4 : Agriculture, forestry and other land use [online]. To be found at <<http://www.ipcc-nggip.iges.or.jp/public/2006gl/vol4.html>> [quoted 25.01.2013]
- Ivánics J, Kovács AZ (1999) Zusammensetzung des Kolostrums bei einigen Fleischrinderrassen in Abhängigkeit von der nach dem Abkalben vergangenen Zeit. *Arch Tierz* 42:17-32
- Janssen H (2006) Untersuchungen zum Einfluß unterschiedlicher Fütterungsintensitäten auf das Körperwachstum, die Futteraufnahme, die Körperzusammensetzung sowie den Stoff- und Energieansatz von weiblichen Kälbern und Aufzuchtrindern. Göttingen : Cuvillier, 171 p
- Kirchgeßner M, Roth FX, Schwarz FJ, Stangl G (2008) Tierernährung : Leitfaden für Studium, Beratung und Praxis. Frankfurt a M : DLG-Verl, 635 p
- Kirchgeßner M, Windisch W, Müller HL (1995) Nutritional factors for the quantification of methane production. In: von Engelhardt W, Leonhard-Marek S, Breves G, Gieseke D (eds) *Ruminant physiology : digestion, metabolism, growth and reproduction : proceedings of the eighth International Symposium on Ruminant Physiology*. Stuttgart : Enke, pp 333-348
- KTBL-Kuratorium für Technik und Bauwesen in der Landwirtschaft (2006) Betriebsplanung Landwirtschaft 2006/07. Darmstadt : KTBL, 672 p
- KTBL-Kuratorium für Technik und Bauwesen in der Landwirtschaft (2010) Betriebsplanung Landwirtschaft 2010/11. Darmstadt : KTBL, 784 p
- Kurihara M, Nishida T, Purnomoadi A, Shibata M, Terada F (2002) The prediction of methane conversion rate from dietary factors. In: Takahashi J, Young BA (eds) *Greenhouse gases and animal agriculture : proceedings of the 1<sup>st</sup> International Conference on Greenhouse Gases and Animal Agriculture*, Obihiro, Japan, 7-11 November 2001. Amsterdam : Elsevier, pp 171-174
- Lockyer R (1997) Methane emissions from grazing sheep and calves. *Agric Ecosyst Environ* 66:11-18
- Meyer U (2005) Fütterung von Kälbern und Jungrindern. *Landbauforsch Völknerode SH* 289:127-136
- Morill KM, Conrad E, Lago A, Campbell J, Quigley J, Tyler H (2012) Nationwide evaluation of quality and composition of colostrum on dairy farms in the United States. *J Dairy Sci* 95:3997-4005
- Pennstate College of Agricultural Sciences (2011) Calf rumen images [online]. To be found at <http://www.das.psu.edu/research-extension/dairy/nutrition/calves/rumen> [quoted 25.01.2013]
- Robelin J, Chilliard Y (1989) Short-term and long-term effects of early nutritional deprivation on adipose tissue growth and metabolism in calves. *J Dairy Sci* 72:505-513
- Schönhusen U, Zitnan R, Kuhla S, Jentsch W, Derno M, Voigt J (2003) Effects of protozoa on methane production in rumen and hindgut of calves around time of weaning. *Arch Anim Nutr* 57:279-295
- Stamer E, Hafez S, Junge W, Kalm E (2004) Genetische Parameter für das Geburts- und Absetzgewicht von schwarzbunten weiblichen Kälbern. *Züchtungskunde* 76:188-195
- Teepker G (2007) Praktische Aspekte der Fütterung einschließlich Haltung und Aufzucht von Kälbern und Mastbullen. *Landbauforsch SH* 313:157-173
- Webb J (2001) Estimating the potential for ammonia emissions from livestock excreta and manures. *Environ Pollut* 111:395-406
- Weiß J, Pabst W, Strack KE, Granz S (2005) Tierproduktion. Stuttgart : Parey, 579 p