

- EL-KASSABY, Y. A., K. RITLAND, A. M. K. FASHLER and W. J. B. DEVITT (1988): The role of reproductive phenology upon the mating system of a Douglas-fir seed orchard. *Silvae Genetica* **37** (2): 76–82.
- EL-KASSABY, Y. A., M. K. FASHLER and O. SZIKLAI (1984): Reproductive phenology and its impact on genetically improved seed production in a Douglas-fir seed orchard. *Silvae Genetica* **33**: 120–125.
- ERICKSON, V. J. and W. T. ADAMS (1989): Mating success in a costal Douglas-fir seed orchard as affected by distance and floral phenology. *Canadian Journal of Forest Research* **19**: 1248–1255
- ERIKSSON, G., A. JONSSON and D. LINDGREN (1973): Flowering in a clonal trial of *Picea abies* Karst. *Stud. For. Suc.*, **110**: 5–45.
- GRIFFIN, A. R. (1984): Clonal variation in Radiata pine seed orchards. II. Flowering phenology. *Australian Forest Research* **14**: 271–281.
- JONSSON, A., I. EKBERG and G. ERIKSSON (1976): Flowering in a seed orchard of *Pinus sylvestris* L. *Stud. For. Suec.* 135pp.
- LI, W., X. WANG and Y. LI (2011): Stability in and Correlation between Factors Influencing Genetic Quality of Seed Lots in Seed Orchard of *Pinus tabulaeformis* Carr. over a 12-Year Span. *PLoS ONE* **6**(8): e23544. doi:10.1371/journal.pone.0023544.
- LI, Y. and X. H. SHEN (1994): Re-selection of clones and seed orchard roguing of *Pinus tabulaeformis* Carr. Asia-Pacific Symposium on Forest Genetics Improvement, IUFRO, Beijing.
- LI, Y., X. R. WANG, W. LI, X. H. SHEN and J. R. XU (2010): Flowering synchronization stabilities of clones among plant ages in a seed orchard of *Pinus tabulaeformis*. *Journal of Beijing Forestry University* **32** (5): 88–93.
- MATZIRIS, D. (1993): Variation in cone production in a clonal seed orchard of black pine. *Silvae Genetica* **42** (2–3): 136–141.
- MATZIRIS, D. I. (1994): Genetic variation in the phenology of flowering in Black pine. *Silvae Genetica* **43** (5–6): 321–328.
- O'REILLY, C., W. H. PARKER and J. E. BARKER (1982): Effect of pollination period and strobili number on random mating in a clonal seed orchard of *Picea mariana*. *Silvae Genetica* **31** (2–3): 90–94.
- SARVAS, R. (1968): Investigations on the flowering and seed crop in *Picea abies*. *Comm. Inst. For. Fenn.* **67** (5): 5–84.
- SAVOLAINEN, O., K. KÄRKKÄINEN, A. HARJU, T. NIKANNEN and M. RUSANEN (1993): Fertility variation in *Pinus sylvestris*: a test of sexual allocation theory. *American Journal of Botany* **80**: 1016–1020.
- SCHMIDTLING, R. C. (1983): Genetic variation in fruitfulness in a Loblolly pine (*Pinus taeda* L.) seed orchard. *Silvae Genetica* **32** (3–4): 76–80.
- SHEN, X. H., Y. LI and X. R. WANG (1985): Study on flowering habit of *Pinus tabulaeformis* Carr in the Seed orchard located in Xingcheng county, Liaoning province. *Journal of Beijing Forestry University* **3**: 1–13.
- WANG, S. S., X. H. SHEN, L. H. YAO, Y. LI, J. G. LI and A. D. LI (1985): A comparative study on growth patterns of *Pinus tabulaeformis* clones in Xingcheng seed orchard, Liaoning province. *Journal of Beijing Forestry College* **4**: 72–83.
- ZAS, R., E. MERLO and J. FERNÁNDEZ-LÓPEZ (2003): SYNCHRO: A SAS Program for Analysing the Floral Phenological Synchronisation in Seed Orchards. *Silvae Genetica* **52** (5–6): 212–215.
- ZHANG D. M., H. X. ZHANG, X. H. SHEN and Y. LI (2004): Study on temporal and spatial change of the mating system in a seed orchard of *Pinus tabulaeformis* Carr. *Scientia Silvae Sinicae* **40** (1): 70–77.
- ZHANG, H. X., G. M. LU and Z. H. LI (1995): An index of phenology overlap in flowering for *Pinus tabulaeformis* Carr. Seed orchard located in Lushi Henan. *Hebei Journal of Forestry and Orchard Research* **10** (3): 199–205.

## Optimisation of a multiplex PCR assay of nuclear microsatellite markers for population genetics and clone identification in *Robinia pseudoacacia* L.

By H. LIESEBACH<sup>1)</sup> and E. EWALD<sup>2)</sup>

(Received 11<sup>th</sup> October 2011)

### Abstract

Black locust (*Robinia pseudoacacia* L.) is a tree species native to North America. The multipurpose tree is cultivated worldwide, but causes problems due to its partially invasive character. The application of nuclear

microsatellite loci has many aims in population genetic studies. Here we introduce a very cost-effective method for combining the information of 14 nuclear microsatellite loci into two multiplex PCR sets as a contribution to greater standardisation and more comparable results.

Combined non-exclusion probabilities for clone identification using example populations are estimated at between  $1.37 \times 10^{-5}$  and  $1.67 \times 10^{-11}$ , and for paternity analysis for  $1.59 \times 10^{-4}$ . The detected weak linkage between some microsatellite loci is not considered to be a substantial restriction to the reliability of the set of

<sup>1)</sup> Thünen Institute of Forest Genetics, Sieker Landstraße 2, D-22927 Grosshansdorf. Phone ++49-4102-696158, Fax ++49-4102-696-200. E-Mail: [heike.liesebach@vti.bund.de](mailto:heike.liesebach@vti.bund.de)

<sup>2)</sup> Thünen Institute of Forest Genetics, Eberswalder Chaussee 3A, D-15377 Waldsiedersdorf.

markers in providing an appropriate method for fingerprinting and parentage analysis.

**Key words:** Black locust; SSRs; multiplexing; population genetics; clone identification; paternity analysis; linkage.

## Introduction

The tree species black locust (*Robinia pseudoacacia* L., Fabaceae) is native to the south-eastern part of the United States and occurs in mountainous regions as an early colonizer of disturbed areas. Black locust has vegetative propagation ability via root suckers and can perform clonal structures in native stands, as was detected by studies with isozyme markers (MCCAIG et al., 1993; CHANG et al., 1998). Black locust flowers are pollinated by insects, and seeds are dispersed by the wind. First mating system observations to describe the generative propagation were carried out by SURLLES et al. (1990). Based on a multilocus isozyme study, they mainly found outcrossing mating, but with remarkable shares of inbreeding (average outcrossing 0.83, range 0.46–1.00).

Black locust, as a „multi-purpose tree,“ is of increasing importance for forestry in many countries. It is cultivated for its durable timber, and for biomass and honey production. Numerous breeding efforts in many countries have been undertaken on the family and clonal level (SCHRÖCK, 1953; KERESZTESI, 1983; MEBRAHTU and HANOVER, 1989; BONGARTEN et al., 1992; DINI-PAPANASTASI, 2008), including the development of several methods for vegetative propagation (NAUJOKS et al., 1999; RÉDEI et al., 2002). As an early colonizer, black locust is suitable for recultivation and slope plantings, but this disposition is also responsible for its partially invasive character. Black locust can become an invasive species in some regions, as ascertained, for example, in Germany (BÖCKER and DIRK, 1998; BÖHMER et al., 2001) and Japan (JUNG et al., 2009). In these countries, clonal structures have been observed in artificial stands (HERTEL and SCHNECK, 2003; JUNG et al., 2009).

The development of highly variable microsatellite markers by LIAN and HOGETSU (2002, seven markers), LIAN et al. (2004, three markers) and MISHIMA et al. (2009, eleven markers) followed the increasing interest in more detailed knowledge of genetic structures in natural and artificial, including invasive, black locust stands and their mating systems, as well as in breeding material. Two studies were published recently describing the application of only four microsatellite loci to recognize clonal structures in artificial populations in Japan (JUNG et al., 2009; KUROKOCHI et al., 2010).

GUICHOUX et al. (2011) reviewed the current trends in microsatellite genotyping and emphasized the need for standardised protocols to enable more comparable results in joint projects. This issue is also under discussion for other species being bred and cultivated, i.e. poplar (RATHMACHER et al., 2009), beech (LEFÈVRE et al., 2011), tomato and wheat (VOSMAN et al., 2001) or Pacific salmon (MORAN et al., 2006; ELLIS et al., 2011).

In this study, we developed a cost-effective multiplex PCR method with 14 loci for further population genetic studies, including mating system analysis and identifi-

cation purposes, and we present first data on linkage and variation levels in a larger number of samples to give a dimension of non-exclusion probabilities for mating system studies. The presented multiplex sets were applied in two studies (LIESEBACH, 2012; LIESEBACH and NAUJOKS, 2012).

## Materials and Methods

### Plant material

The plant material originates from several samples, mainly from artificial stands in Germany. Among them are semi-natural populations which have not been managed for a long time; selected clones; material from actual breeding programs; open pollinated offspring from single clones, and samples from six planted stands from several seed sources located in Germany/Brandenburg. In addition to *Robinia pseudoacacia*, a few samples of other *Robinia* species were tested: an open pollinated offspring family from *R. neomexicana* A. Gray (37 individuals) and the clone ‘Casque Rouge’ (*R.* × *margarettiae* Ashe, *R. hispida* L. × *R. pseudoacacia*). The previous total sample size amounts to approximately 1300 individuals.

### Microsatellite genotyping

The isolation of total DNA from fresh or frozen leaves or from seeds followed a modified CTAB protocol (DUMOLIN et al., 1995). All 21 available loci were tested with the exception of Rops15, which was described as highly somatic instable (LIAN et al., 2004), and RP102, RP211 and RP165, which were characterized by a low variation and a significant departure from Hardy-Weinberg equilibrium (MISHIMA et al., 2009).

PCRs for the remaining 17 microsatellite loci (Table 1) were carried out with fluorescent dye-labeled forward primers (delivered by biomers.net, [www.biomers.net](http://www.biomers.net)) using the “Multiplex PCR Kit” from Qiagen in accordance with the manufacturer’s instructions. PCRs were carried out in 15 µl reaction volume with 10–50 ng template DNA in the Multiplex PCR Master Mix containing HotStarTaq DNA Polymerase, buffer, dNTP mix and a final concentration of 3 mM MgCl<sub>2</sub>. All primers were added with equal concentrations of 0.2 µM. PCRs were carried out in a Biometra TGradient and a Biometra UNOII thermocycler (Göttingen, Germany) with the following temperature profiles: activation step of 15 minutes at 95°C, 28 cycles (denaturation of 30 sec at 94°C, annealing 90 sec, and extension for 60 sec at 72°C) and a final extension of 30 min at 60°C. After optimisation, eight loci were analysed in Multiplex Set 1 at an annealing temperature of 56°C, six loci were analysed in Multiplex Set 2 at an annealing temperature of 63° (Table 1) and the three loci Rops09, Rops10 and Rops18 were excluded (see below). Amplification products were detected with a Beckman Coulter CEQ 8000 capillary sequencer.

### Data analysis

Population genetic parameters such as number of alleles per locus (*A/L*) and observed heterozygosity (*H<sub>o</sub>*) as

well as combined non-exclusion probabilities for the identification of unrelated individuals, sib identity or the identification of the second parent when the first parent is known, were calculated with the software package CERVUS (MARSHALL et al., 1998; KALINOWSKI et al., 2007). The effective number of alleles  $N_e$  was estimated by the reciprocal of expected homozygosity.

Homozygous genotypes of null alleles were assumed in the case of repeated missing amplification products in samples with other detectable loci. A further indication for null alleles in heterozygote genotypes is mismatching between mother and offspring individuals when mother and offspring exhibit different alleles in apparent homozygous genotypes. The software package Micro-

Table 1. – Overview of nuclear microsatellite loci applied on *Robinia* samples and their variation parameters.

Locus	Reference for primer development	Motif	PCR set	Dye used to label the forward primer	Range (bp)	Number of alleles in <i>R. pseudo-acacia</i>	Observed heterozygosity	Additional alleles in <i>R. neomexicana</i> and <i>R. hispida</i>
Rops04	(Lian and Hogetsu, 2002)	AC	1	BMN-6	107-112	4 <sup>n</sup>	0.3153	1
Rops05	(Lian and Hogetsu, 2002)	AC	1	DY-751	115-156	14	0.8312	2
Rops06	(Lian and Hogetsu, 2002)	GT	1	Cy5	117-147	10	0.6841	
RP106	(Mishima et al., 2009)	GT	1	BMN-6	128-138	4	0.6321	1
RP01B	(Mishima et al., 2009)	CT	1	Cy5	155-175	9	0.8165	1
Rops10	(Lian and Hogetsu, 2002)	T/AAT	1, test only	BMN-6	181-188	6 <sup>n</sup>	0.1659	
RP150	(Mishima et al., 2009)	TC	1	BMN-6	178-202	12	0.8187	
Rops08	(Lian and Hogetsu, 2002)	CA	1	Cy5	193-207	7	0.6741	
Rops16	(Lian et al., 2004)	CT	1	DY-751	194-220	13	0.8627	1
Rops09	(Lian and Hogetsu, 2002)	TA/A	2, test only	BMN-6	77-127	10 <sup>n</sup>	0.0568	
RP035	(Mishima et al., 2009)	TC	2	BMN-6	75- 97	10 <sup>n</sup>	0.6805	
RP032	(Mishima et al., 2009)	TG	2	DY-751	91-119	11 <sup>n</sup>	0.2913	
Rops02	(Lian and Hogetsu, 2002)	AC/AT	2	Cy5	106-142	14 <sup>n</sup>	0.7866	2
RP109	(Mishima et al., 2009)	AG	2	BMN-6	123-145	11	0.7292	2
RP200	(Mishima et al., 2009)	AG	2	DY-751	136-182	19	0.7814	1
Rops18	(Lian et al., 2004)	AC	2, test only	DY-751	137-221	10	0.6837	
RP206	(Mishima et al., 2009)	GT	2	Cy5	205-231	13 <sup>n</sup>	0.5799	1

<sup>n</sup> assumed null allele because of missing amplification products or mismatches between mother and offspring.

Checker (VAN OOSTERHOUT et al., 2004) was used to estimate null allele frequencies based on Hardy-Weinberg equilibrium in some example populations. One of the implemented methods (BROOKFIELD, 1996) considers missing values as true homozygote genotypes of the null allele.

Linkage analyses were carried out for some two-locus combinations derived from available mother-offspring families. Haploid maternal genotypic data were generated from the offspring by subtracting the paternal alleles. Data were omitted in case of identity of maternal and offspring heterozygote genotype since the maternal allele cannot be determined. Recombination frequencies  $r$  with standard deviations were calculated as

$$r = \frac{R}{N} \pm \sqrt{\frac{\frac{R}{N} * (1 - \frac{R}{N})}{N}}$$

where  $N$  is the total sample size of two-locus combinations and  $R$  is the number of recombinants. Kosambi's mapping function was used to calculate map distances as

$$d = \frac{1}{4} \ln \frac{1 + 2r}{1 - 2r}$$

The classical LOD scores of linkage analysis was calculated as

$$\text{LOD} = \log_{10} \frac{(1 - \frac{R}{N})^{n-\kappa} * (\frac{R}{N})^{\kappa}}{0.5^N}$$

as the decimal logarithm of the likelihood ratio of linkage between the two loci and of independent segregation. The common threshold is  $\text{LOD} \geq 3.0$  for significant linkage as was suggested by MORTON (1955), cited in GERBER and RODOLPHE (1994).

## Results and Discussion

### Variation

Initially, 17 microsatellite loci were analysed. After the first approximately 200 samples, three loci were excluded from further analysis to concentrate a maximum of information in a minimum of PCRs. Locus Rops09 was omitted because of its very low variation and a putative null allele (observed heterozygosity 0.057). Locus Rops10 has a rather mononucleotide pattern and was omitted because of a putative highly frequent null allele. About 70% of individuals did not reveal a peak at this locus. Locus Rops18 was excluded because of its wide range from 137 to 221 bp with no alleles between 152 and 209 bp, perhaps caused by a large indel overlaying the dinucleotide pattern. A tendency to allele dropout is suspected because large alleles do not amplify as efficiently as small alleles. Some extreme differences in peak height were observed in heterozygous genotypes combining alleles shorter than 152 bp and longer than 209 bp. The remaining 14 loci were combined in two multiplex PCR sets with 8 and 6 loci, respectively (Table 1).

All loci more or less fit to a dinucleotide base-pair periodicity with the exception of locus Rops05 with a

dinucleotide repeat that exhibits an average of 2.2 nucleotides distance between adjacent alleles. This so-called size shift exists between the observed electrophoretic size and the expected repeat unit difference. Similar deviations from the expected periodicity were observed in humans (AMOS et al., 2007), in poplar (LIESEBACH et al., 2010) and salmon (ELLIS et al., 2011). Alleles of *R. neomexicana* also match to the *R. pseudoacacia* ladders, as was observed in an offspring family. However, we detected alleles apart from the ladder at five loci in the clone 'Casque Rouge', obviously originating from the *R. hispida* parent.

Even though these 14 loci are suitable for clone identification, one locus has to be excluded for population genetic studies, especially mating system and paternity analysis. Locus RP150 shows an indication for duplication. It reveals reproducible genotype patterns with 1 to 4 alleles per sample and partially shows behaviour like a tetraploid locus with a dosage effect, i.e., two small peaks and one large peak.

### Linkage

Multilocus data evaluations, like the calculation of several non-exclusion probabilities, assume that the loci segregate independently. However, no serious bias is expected in multilocus calculations when loci are not so tightly linked (MARSHALL et al., 1998). Mapping distances of 10 cM or more are considered as an independent association of markers for fingerprinting and parentage analysis purposes (SLAVOV et al., 2004).

Here we present first results of linkage analysis in black locust in offspring families from two openly pollinated trees A and B. Forty five two-locus combinations could be tested from 10 heterozygote loci for each of the individuals A and B. Table 2 shows five significant linkages at the LOD score > 3 criterion. The suggested threshold value of 3 is very conservative, as was discussed by GERBER and RODOLPHE (1994). Yet further two-locus combinations in this study have LOD scores below 1.6 and have to be considered as independent loci. Two pairs of linked loci Rops05 – RP035 and RP109 – RP200 were observed in both trees A and B, whereas RP106 – RP01B in tree B (Table 2) seems to be unlinked in tree A (LOD = 0.0098, recombination frequency 0.489).

### Estimation of non-exclusion probabilities

Generally, non-exclusion probabilities should be as small as possible. This could be achieved by a high number of loci and a high level of variation. Nevertheless, it could be advisable to exclude single loci for paternity analysis in the case of high frequencies of null alleles in certain populations. Often they cannot contribute information because of missing amplification products, or their apparent homozygous genotype might be a true heterozygote with the null allele. In the six planted stands of black locust described here, maximum observed frequencies of null alleles per population are 18% for locus Rops04; 23% for RP032, and 29% for RP206. However, such loci can be used as additional information in paternity analysis to exclude candidate fathers in the case of more than one possible candidate.

Table 2. – Linkage of nuclear microsatellite loci in *R. pseudoacacia*.

Locus1	Locus2	Tree ID	Sample size	Segregation ratio in offspring	LOD score	Recombination frequency $r \pm \text{st.dev}$	Map distance $d$ (cM)
Rops05	RP035	A	173	88 : 14 : 13 : 58	19.539	0.156 ± 0.028	16.15
Rops05	RP035	B	63	21 : 1 : 8 : 33	7.744	0.143 ± 0.044	14.69
RP109	RP200	A	159	101 : 14 : 10 : 34	18.562	0.151 ± 0.028	15.58
RP109	RP200	B	69	36 : 4 : 0 : 29	14.138	0.058 ± 0.028	5.82
RP106	RP01B	B	44	6 : 10 : 25 : 3	3.564	0.205 ± 0.061	21.73

Table 3. – Examples of variation parameters and combined non-exclusion probabilities for six artificial *R. pseudoacacia* stands based on evaluation of 13 highly variable microsatellite loci.

Population	Number of individuals	Allele per Locus (A/L)	Effective number of alleles (Ae)	Observed heterozygosity (Ho)	Combined non-exclusion probability (identity)	Combined non-exclusion probability (sib identity)	Combined non-exclusion probability (second parent)
1	74	9.62	5.2164	0.7128	1.26E-15	3.87E-6	5.58E-6
2	74	6.23	3.1791	0.7155	1.00E-10	5.72E-5	9.18E-4
3	69	9.31	5.2753	0.7106	2.06E-15	4.67E-6	6.40E-6
4	74	8.15	4.9288	0.6912	6.69E-15	5.84E-6	1.15E-5
5	72	9.15	5.0532	0.6748	5.06E-15	6.18E-6	9.52E-6
6	72	9.15	5.3216	0.7325	1.21E-15	4.21E-6	5.21E-6
Average	72.5	8.60	4.8291	0.7062	1.67E-11	1.37E-5	1.59E-4

Some examples for variation parameters and combined non-exclusion probabilities based on 13 microsatellite loci (locus RP150 was excluded because of duplication) were given for six populations to demonstrate the power of the presented set of highly variable microsatellite loci in black locust (Table 3). Calculations were carried out under the condition of unlinked loci. Taking into consideration the observed linkage, the combined non-exclusion probabilities might be slightly underestimated.

A realistic probability for identifying clones originating from vegetative propagation in populations or in breeding materials by their identical multilocus genotypes might be assumed between the estimates for identity by chance for unrelated individuals (on average 1.67E-11) and for sibs (on average 1.37E-05). This implies a highly reliable identification of clones including possible full sibs. The application of the set of

microsatellite markers for parentage analysis is not restricted by linkage despite inexact non-exclusion probabilities. It allows a reliable identification of the pollinator among the candidate fathers or the exclusion of all candidates in natural or semi-natural populations with clonal structures or in seed orchards.

#### Acknowledgements

The authors thank VOLKER SCHNECK, GISELA NAUJOKS, JAN ENGEL and DIRK KNOCHÉ for their assistance in the collection of *R. pseudoacacia* plant material and Roland Graeff for providing the *R. neomexicana* collection. Partial funding was provided by the Fachagentur Nachwachsende Rohstoffe e.V. (FNR) of the Federal Ministry of Food, Agriculture and Consumer Protection in Germany. The authors also thank two anonymous reviewers for their constructive contributions to improve the manuscript.

## References

- AMOS, W., J. I. HOFFMAN, A. FRODSHAM, L. ZHANG, S. BEST and A. V. S. HILL (2007): Automated binning of microsatellite alleles: problems and solutions. *Molecular Ecology Notes* **7**: 10–14.
- BÖCKER, R. and M. DIRK (1998): Distribution and Spreading of Alien Trees and Shrubs in South Western Germany and Contributions to Germination Biology. pp. 285–297. *In*: Plant Invasions: Ecological Mechanisms and Human Responses, edited by U. STARFINGER, K. EDWARDS, I. KOWARIK and M. WILLIAMSON, Backhuys Publishers, Leiden, The Netherlands.
- BÖHMER, H. J., T. HEGER and L. TREPL (2001): *Robinia pseudoacacia* L., Black locust. pp. 64–75. *In*: Fallstudien zu gebietsfremden Arten in Deutschland – Case Studies on Alien Species in Germany, Umweltbundesamt (Federal Environment Agency), Berlin.
- BONGARTEN, B. C., D. A. HUBER and D. K. APSLEY (1992): Environmental and genetic influences on short-rotation biomass production of black locust (*Robinia pseudoacacia* L.) in the Georgia Piedmont. *Forest Ecology and Management* **55**: 315–331.
- BROOKFIELD, J. F. Y. (1996): A simple new method for estimating null allele frequency from heterozygote deficiency. *Molecular Ecology* **5**: 453–455.
- CHANG, C.-S., B. BONGARTEN and J. HAMRICK (1998): Genetic Structure of Natural Populations of Black Locust (*Robinia pseudoacacia* L.) at Coweeta, North Carolina. *Journal of Plant Research* **111**: 17–24.
- DINI-PAPANASTASI, O. (2008): Effects of clonal selection on biomass production and quality in *Robinia pseudoacacia* var. *monophylla* Carr. *Forest Ecology and Management* **256**: 849–854.
- DUMOLIN, S., B. DEMESURE and R. J. PETIT (1995): Inheritance of chloroplast and mitochondrial genomes in pedunculate oak investigated with an efficient PCR method. *TAG Theoretical and Applied Genetics* **91**: 1253–1256.
- ELLIS, J. S., J. GILBEY, A. ARMSTRONG, T. BALSTAD, E. CAUWELIER, C. CHERBONNEL, S. CONSUEGRA, J. COUGHLAN, T. F. CROSS, W. CROZIER, E. DILLANE, D. ENSING, C. G. D. LEÁNIZ, E. GARCÍA-VÁZQUEZ, A. M. GRIFFITHS, K. HINDAR, S. HJORLEIFSDOTTIR, D. KNOX, G. MACHADO-SCHIAFFINO, P. MCGINNITY, D. MELDRUP, E. E. NIELSEN, K. OLAFSSON, C. R. PRIMMER, P. PRODOHL, L. STRADMEYER, J.-P. VÁHÁ, E. VERSPOOR, V. WENNEVIK and J. R. STEVENS (2011): Microsatellite standardization and evaluation of genotyping error in a large multi-partner research programme for conservation of Atlantic salmon (*Salmo salar* L.). *Genetica* **139**: 353–367.
- GERBER, S. and F. RODOLPHE (1994): Estimation and test for linkage between markers: a comparison of lod score and  $X^2$  test in a linkage study of maritime pine (*Pinus pinaster* Ait.). *TAG Theoretical and Applied Genetics* **88**: 293–297.
- GUICHOUX, E., L. LAGACHE, S. WAGNER, P. CHAUMEIL, P. LÉGER, O. LEPAIS, C. LEPOITTEVIN, T. MALAUSA, E. REVARDEL, F. SALIN and R. J. PETIT (2011): Current trends in microsatellite genotyping. *Molecular Ecology Resources* **11**: 591–611.
- HERTEL, H. and V. SCHNECK (2003): Untersuchungen zur genetischen Struktur eines Robinienbestandes (*Robinia pseudoacacia* L.) in Brandenburg. pp. 257–263. *In*: Bedrohung der biologischen Vielfalt durch invasive gebietsfremde Arten – Erfassung, Monitoring und Risikoanalyse, edited by M. Welling, Landwirtschaftsverlag GmbH, Münster.
- JUNG, S.-C., N. MATSUSHITA, B.-Y. WU, N. KONDO, A. SHIRAISHI and T. HOGETSU (2009): Reproduction of a *Robinia pseudoacacia* population in a coastal *Pinus thunbergii* windbreak along the Kujukurihama Coast, Japan. *Journal of Forest Research* **14**: 101–110.
- KALINOWSKI, S. T., M. L. TAPER and T. C. MARSHALL (2007): Revising how the computer program CERVUS accommodates genotyping error increases success in paternity assignment. *Molecular Ecology* **16**: 1099–1106.
- KERESZTESI, B. (1983): Breeding and cultivation of black locust, *Robinia pseudoacacia*, in Hungary. *Forest Ecology and Management* **9**: 217–244.
- KUROKOCHI, H., K. TOYAMA and T. HOGETSU (2010): Regeneration of *Robinia pseudoacacia* riparian forests after clear-cutting along the Chikumagawa River in Japan. *Plant Ecology* **210**: 31–41.
- LEFÈVRE, S., S. WAGNER, R. J. PETIT and G. DE LAFONTAINE (2011): Multiplexed microsatellite markers for genetic studies of beech. *Molecular Ecology Resources* doi: 10.1111/j.1755-0998.2011.03094.x.
- LIAN, C. and T. HOGETSU (2002): Development of microsatellite markers in black locust (*Robinia pseudoacacia*) using a dual-suppression-PCR technique. *Molecular Ecology Notes* **2**: 211–213.
- LIAN, C., R. OISHI, N. MIYASHITA and T. HOGETSU (2004): High somatic instability of a microsatellite locus in a clonal tree, *Robinia pseudoacacia* TAG Theoretical and Applied Genetics **108**: 836–841.
- LIESEBACH, H. (2012): Genetische Charakterisierung von Robinienbeständen (*Robinia pseudoacacia* L.) in Deutschland mit nuklearen Mikrosatelliten-Markern: Erkenntnisse zu ihrer Bestandesbegründung (Genetic characterisation of black locust stands (*Robinia pseudoacacia* L.) in Germany with nuclear microsatellite markers: background on population history). Beiträge aus der Nordwestdeutschen Forstlichen Versuchsanstalt, in press.
- LIESEBACH, H. and G. NAUJOKS (2012): Klonidentifizierung bei Zuchtmaterial der Robinie (*Robinia pseudoacacia* L.) mit nuklearen Mikrosatellitenmarkern (Clone identification of breeding material from black locust (*Robinia pseudoacacia* L.) from nuclear microsatellite markers). Beiträge aus der Nordwestdeutschen Forstlichen Versuchsanstalt, Volume **8**, 267–274.
- LIESEBACH, H., V. SCHNECK and E. EWALD (2010): Clonal fingerprinting in the genus *Populus* L. by nuclear microsatellite loci regarding differences between sections, species and hybrids. *Tree Genetics & Genomes* **6**: 259–269.
- MARSHALL, T. C., J. SLATE, L. E. B. KRUIK and J. M. PEMBERTON (1998): Statistical confidence for likelihood-based paternity inference in natural populations. *Molecular Ecology* **7**: 639–655.
- MCCAIG, B. C., J. L. HAMRICK and B. L. HAINES (1993): Clonal structure of *Robinia pseudoacacia* (black locust) in the Southern Appalachians. *Bulletin of the Ecological Society of America* **74**: 350.
- MEBRAHTU, T. and J. W. HANOVER (1989): Heritability and expected gain estimates for traits of Black locust in Michigan. *Silvae Genetica* **38**: 125–130.
- MISHIMA, K., T. HIRAO, S. URANO, A. WATANABE and K. TAKATA (2009): Isolation and characterization of microsatellite markers from *Robinia pseudoacacia* L. *Molecular Ecology Resources* **9**: 850–852.
- MORAN, P., D. J. TEEL, E. S. LAHOOD, J. DRAKE and S. KALINOWSKI (2006): Standardising multi-laboratory microsatellite data in Pacific salmon: an historical view of the future. *Ecology of Freshwater Fish* **15**: 597–605.

- NAUJOKS, G., I. ZASPEL and U. BEHRENDT (1999): Microorganisms acting in tissue cultures of black locust (*Robinia pseudoacacia* L.). *Acta Horticulturae* **530**: 129–135.
- RATHMACHER, G., M. NIGGEMANN, H. WYPUKOL, K. GEBHARDT, B. ZIEGENHAGEN and R. BIALOZYT (2009): Allelic ladders and reference genotypes for a rigorous standardization of poplar microsatellite data. *Trees – Structure and Function* **23**: 573–583.
- RÉDEI, K., Z. OSVÁTH-BUJTÁS and I. BALLA (2002): Clonal approaches to growing black locust (*Robinia pseudoacacia*) in Hungary: a review. *Forestry* **75**: 547–552.
- SCHRÖCK, O. (1953): Beitrag zur Züchtung der Robinie (*Robinia pseudoacacia*). *Der Züchter* **23**: 266–272.
- SLAVOV, G. T., G. T. HOWE, I. A. YAKOVLEV, K. J. EDWARDS, K. V. KRUTOVSKY, G. A. TUSKAN, J. E. CARLSON, S. H. STRAUSS and W. T. ADAMS (2004): Highly variable SSR markers in Douglas-fir: Mendelian inheritance and map locations. *TAG Theoretical and Applied Genetics* **108**: 873–880.
- SURLES, S. E., J. L. HAMRICK and B. BONGARTEN (1990): Mating systems in open-pollinated families of black locust (*Robinia pseudoacacia*). *Silvae Genetica* **39**: 35–40.
- Van Oosterhout, C., W. F. HUTCHINSON, D. P. M. WILLS and P. SHIPLEY (2004): MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes* **4**: 535–538.
- VOSMAN, B., R. COOKE, M. GANAL, R. PEETERS, P. ISAAC and G. BREDEMEIJER (2001): Standardization and application of microsatellite markers for variety identification on tomato and wheat. *Acta Horticulturae* 546.

## Microarray Analysis of Gene Expression in Triploid Black Poplar

By BAOQUAN HU<sup>1</sup>, BIN WANG<sup>1</sup>, CHUNGUO WANG<sup>1</sup>, WENQIN SONG<sup>1</sup> and CHENGBIN CHEN<sup>1,\*</sup>

(Received 19<sup>th</sup> November 2011)

### Abstract

Triploidy is a widespread phenomenon in cultivated and natural breeding plants and it can confer some growth advantages. Here, we analyzed genome-wide gene expression in triploid *Populus euramericana* (black poplar) using the Affymetrix poplar microarray to detect any possible correlation between triploid vigor and a unique gene expression profile. Among the 38,400 transcripts that were detected in triploid poplar, 1,564 and 2,015 genes were up- or downregulated, respectively, compared with the diploid. The majority of the upregulated genes in the triploid were associated with carbon and nitrogen metabolism, especially lignin and secondary metabolism. Other genes upregulated in the triploid included genes involved in sugar transport, and brassinosteroid (BR) and auxin metabolism. Downregulated genes were mostly related to the assembly and biosynthesis of ribosomes and the nucleosome macromolecular complex. The results suggested that BR and auxin levels were crucial in controlling sugar transport, photosynthesis and cell wall biosynthesis. Downregulated genes were associated with chromatin regulation in the triploid. The information from this analysis could provide an insight into the vigor of triploid poplar.

**Key words:** Gene expression; *Populus*; microarray; triploid.

**Abbreviations:** BR, brassinosteroid; cRNA, complementary RNA; ESTs, expressed-sequence tags; GO, gene ontology; IAA,

indole-3-acetic acid; qRT-PCR quantitative real-time polymerase chain reaction; SEA, singular enrichment analysis.

### Introduction

Polyploidy is a widespread phenomenon in plants, and it is also a major mechanism driving speciation. Approximately 60–70% of flowering plants have undergone polyploidization during their ancestry (BLANC and WOLFE, 2004; CUI et al., 2006; MASTERSON, 1994), and it is estimated that 15% of angiosperm speciation has been achieved by polyploidization (RIESEBERG and WILLIS, 2007). However, new polyploids often face reproductive isolation from their progenitors due to the triploid reproductive barrier, which is caused by the production of unreduced gametes (KÖHLER et al., 2010). Therefore, the plants that evolved by polyploidization remain in a triploid state. This speciation process has been defined as the triploid bridge, and is thought to play an important role in plant evolution (YAMAUCHI et al., 2004).

Triploidy is also found in natural plant groups, and these plants can have growth advantages. There exist triploid varieties in various species of poplar, including *Populus tremula*, *Populus alba* (PAULEY, 1949), *Populus balsamifera* and *Populus tremuloides* (VAN BUIJTENEN et al., 1958). Compared with their diploid counterparts, the triploid trees are faster growing, have larger leaves and show greater vigor (LI et al., 2008). Several studies have examined triploid white poplar, which has good growth performance and other desirable properties (ZHANG et

<sup>1</sup>) College of Life Sciences, NanKai University, Tianjin 300071, People's Republic of China.

\*) Corresponding author: CHENGBIN CHEN. Tel.: +86 22 23508241, Fax: +86 22 23508241. E-Mail address: chchbin@126.com