



## RESEARCH PAPER

# Environmental triggers for photosynthetic protein turnover determine the optimal nitrogen distribution and partitioning in the canopy

Yi-Chen Pao\*, Tsu-Wei Chen, Dany Pascal Moualeu-Ngangue and Hartmut Stützel

Institute of Horticultural Production Systems, Leibniz Universität Hannover, Hannover, Germany

\* Correspondence: [pao@gem.uni-hannover.de](mailto:pao@gem.uni-hannover.de)

Received 28 March 2018; Editorial decision 14 August 2018; Accepted 14 August 2018

Editor: Pierre Martre, INRA, France

## Abstract

Plants continually adjust the photosynthetic functions in their leaves to fluctuating light, thereby optimizing the use of photosynthetic nitrogen ( $N_{ph}$ ) at the canopy level. To investigate the complex interplay between external signals during the acclimation processes, a mechanistic model based on the concept of protein turnover (synthesis and degradation) was proposed and parameterized using cucumber grown under nine combinations of nitrogen and light in growth chambers. Integrating this dynamic model into a multi-layer canopy model provided accurate predictions of photosynthetic acclimation of greenhouse cucumber canopies grown under high and low nitrogen supply in combination with day-to-day fluctuations in light at two different levels. This allowed us to quantify the degree of optimality in canopy nitrogen use for maximizing canopy carbon assimilation, which was influenced by  $N_{ph}$  distribution along canopy depth or  $N_{ph}$  partitioning between functional pools. Our analyses suggest that  $N_{ph}$  distribution is close to optimum and  $N_{ph}$  reallocation is more important under low nitrogen.  $N_{ph}$  partitioning is only optimal under a light level similar to the average light intensity during acclimation, meaning that day-to-day light fluctuations inevitably result in suboptimal  $N_{ph}$  partitioning. Our results provide insights into photoacclimation and can be applied to crop model improvement.

**Keywords:** Functional partitioning, light, mechanistic model, nitrogen reallocation, nitrogen supply, optimal, photosynthetic acclimation.

## Introduction

Acclimation of leaf traits to fluctuating environments is a key mechanism to maximize fitness (Walters, 2005; Athanasiou *et al.*, 2010). To maximize canopy carbon gain, dynamic modifications of photosynthetic traits to track heterogeneous light distribution within the canopy are crucial (Retkute *et al.*, 2015), especially for herbaceous species with a continuously leaf-forming nature (Niinemets *et al.*, 2015). One of the most important strategies in photoacclimation is to maintain efficient

utilization of limited resources in the photosynthetic apparatus, e.g. nitrogen, by continuous modifications of (i) between-leaf distribution along the canopy depth and (ii) within-leaf partitioning between photosynthetic functions according to local light availability (Evans, 1989).

Vertical nitrogen distribution in response to light has been intensively studied (Hirose and Werger, 1987; Werger and Hirose, 1991; Anten *et al.*, 1995; Dreccer *et al.*, 2000; Moreau

Abbreviations: aDPI, average daily photosynthetic photon integral during acclimation; Chl, chlorophyll per unit leaf area; DCA, daily canopy carbon assimilation; DPI, daily photosynthetic photon integral; HL, high light; HN, high nitrogen;  $J_{max}$ , maximum electron transport rate; LL, low light; LAI, leaf area index; LN, low nitrogen;  $N_c$ , light harvesting pool of photosynthetic nitrogen;  $N_e$ , electron transport pool of photosynthetic nitrogen;  $N_{ph}$ , photosynthetic nitrogen;  $N_v$ , carboxylation pool of photosynthetic nitrogen; PNUE, photosynthetic nitrogen use efficiency; PPFD, photosynthetic photon flux density;  $V_{cmax}$ , maximum carboxylation rate.

© The Author(s) 2018. Published by Oxford University Press on behalf of the Society for Experimental Biology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

*et al.*, 2012; Hikosaka *et al.*, 2016). Nitrogen distribution was reported to closely follow the light gradient and thus approach its optimum in wheat stands (Dreccer *et al.*, 2000). However, this relationship has not been found in other studies (Moreau *et al.*, 2012; Hikosaka *et al.*, 2016). In fact, many studies demonstrated that nitrogen distribution failed to track the within-canopy light gradient optimally due to a delay in nitrogen reallocation in the lower canopy layer and an underinvestment in the upper layer (Field, 1983; Evans, 1993; Hollinger, 1996; Hirose *et al.*, 1997; Meir *et al.*, 2002; Wright *et al.*, 2006; Hikosaka, 2016). This discrepancy between optimum and reality could be explained by physiological limitations and the cost of nitrogen reallocation (Hikosaka, 2016; Kitao *et al.*, 2018) or might result from incorrect predictions. In some cases (e.g. Hikosaka, 2014; Kitao *et al.*, 2018), the optimal nitrogen distribution that followed the within-canopy light gradient estimated by the Beer-Lambert law was predicted to be extremely high in the upper canopy, which might not be biologically reachable. This could result from the oversimplification of models in three aspects: (i) neglecting the effects of variations in the structural characteristics, e.g. leaf elevation angle (Falster and Westoby, 2003), on light interception of the leaves; (ii) neglecting age-dependent modifications and limitations during leaf development and ageing (Niinemets *et al.*, 2015; Niinemets, 2016); and (iii) assuming a linear relationship between photosynthetic capacity and photosynthetic nitrogen per unit leaf area instead of considering photoacclimation in functional nitrogen partitioning.

Optimizing functional partitioning within the leaf is of great importance because it improves carbon gain by enhancing photosynthetic nitrogen use efficiency (PNUE; Zhu *et al.*, 2010). Photosynthetic rate is determined by the limited rate of ribulose 1,5-bisphosphate (RuBP) carboxylation and RuBP regeneration in the photosynthetic machinery (Farquhar *et al.*, 1980). Besides driving photosynthesis, light also triggers fine adjustments in nitrogen investment between (i) RuBP carboxylation (Rubisco), (ii) RuBP regeneration (electron transport), and (iii) light harvesting functions (Yamori *et al.*, 2010; Trouwborst *et al.*, 2011; Vialet-Chabrand *et al.*, 2017). The capability and significance of photoacclimation in functional nitrogen partitioning were empirically addressed in both light-demanding and shade-tolerant species (Evans, 1993; Hikosaka and Terashima, 1996; Pons and Anten, 2004; Hikosaka, 2005; Trouwborst *et al.*, 2011). Recently, with a modelling approach, it was predicted that a decreasing investment in the light harvesting function can increase canopy PNUE (Song *et al.*, 2017). However, genetic and physiological controls of photoacclimatory processes by environmental triggers are still not described mechanistically.

The degree of acclimation under a given environment is limited by the previous environmental conditions (Walters, 2005; Niinemets *et al.*, 2006) along with continuous age-dependent modifications in physiological traits (Niinemets, 2016). This emphasizes that static models, which do not consider the dynamics of plant growth and environmental fluctuations, may not be sufficiently precise in predicting acclimation behavior. Prieto *et al.* (2012) proposed an empirical model describing the combined effects of leaf age and light on leaf nitrogen economics for a grapevine canopy and demonstrated that the mean daily light integral over the previous 10 d explained 73%

of the variation in nitrogen per unit leaf area. Since environmental acclimation and developmental (genetic control of leaf ageing) acclimation are regulated distinctively (Athanasίου *et al.*, 2010), it is possible to integrate internal (age) and external (environment) triggers into a mechanistic model for better understanding of the developmental and environmental effects on photosynthetic acclimation.

Acclimation processes in leaf functioning are regulated by constant updates of protein content as a result of protein turnover, driven by the concurrent actions of degradation and synthesis (Li *et al.*, 2017). In growing leaves, photosynthetic proteins account for the highest cost in protein turnover (Li *et al.*, 2017). At the expense of energy, protein turnover is necessary for adjusting protein levels in line with external triggers. It was experimentally shown that leaf Rubisco content increased with light (Yamori *et al.*, 2010) and nitrogen supply level (Yamori *et al.*, 2011a) and exhibited an evolution with leaf age that could be interpreted by Rubisco turnover (Suzuki *et al.*, 2001; Ishimaru *et al.*, 2001; Irving and Robinson, 2006). Based on the concept of protein turnover, Thornley (1998) proposed a mechanistic model predicting reasonable dynamics of photosynthetic acclimation at the leaf level. We refined this model to describe the dynamics of different photosynthetic nitrogen pools and to quantify the developmental and environmental effects of light and nitrogen availabilities on leaf acclimation. The optimality of nitrogen distribution and partitioning at the canopy scale was evaluated by integrating this model into a multi-layer model considering the structural characteristics of a cucumber canopy. This aims (i) to test whether the protein turnover can be a mechanistic explanation of the photosynthetic acclimation under dynamic environmental conditions; and (ii) to understand the regulatory mechanism of environmental triggers on the degree of optimality at the canopy level in terms of maximizing PNUE and canopy carbon assimilation, which can be considered as an indicator of the general fitness of the plants.

## Materials and methods

### Modelling the dynamics of photosynthetic protein turnover

Photosynthetic nitrogen ( $N_{\text{ph}}$ , mmol N m<sup>-2</sup>) is defined as biologically active nitrogen in the proteins involved in photosynthetic functions, i.e. carboxylation, electron transport and light harvesting. Leaf  $N_{\text{ph}}$  is calculated as the sum of nitrogen in the carboxylation pool ( $N_{\text{v}}$ ), electron transport pool ( $N_{\text{j}}$ ) and light harvesting pool ( $N_{\text{c}}$ ; Trouwborst *et al.*, 2011):

$$N_{\text{ph}} = N_{\text{v}} + N_{\text{j}} + N_{\text{c}} \quad (1)$$

where  $N_{\text{v}}$  includes only Rubisco and represents the nitrogen investment in carboxylation capacity,  $N_{\text{j}}$  includes the electron transport chain, photosystem II core and Calvin cycle enzymes other than Rubisco, and  $N_{\text{c}}$  includes the photosystem I core and light harvesting complexes I and II. Functional pools  $N_{\text{v}}$ ,  $N_{\text{j}}$ , and  $N_{\text{c}}$  are estimated from the maximum carboxylation rate ( $V_{\text{cmax}}$ ,  $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ ), maximum electron transport ( $J_{\text{max}}$ ,  $\mu\text{mol e}^- \text{ m}^{-2} \text{ s}^{-1}$ ) and leaf chlorophyll ( $\text{Chl}$ , mmol Chl m<sup>-2</sup>), respectively (Buckley *et al.*, 2013):

$$N_{\text{v}} = V_{\text{cmax}} / \chi_{\text{v}} \quad (2a)$$

$$N_{\text{j}} = J_{\text{max}} / \chi_{\text{j}} \quad (2b)$$

$$N_C = (Chl - N_J \times \chi_{Cj}) / \chi_C \quad (2c)$$

where  $\chi_V$  ( $\mu\text{mol CO}_2 \text{ mmol}^{-1} \text{ N s}^{-1}$ ) is the carboxylation capacity per unit Rubisco nitrogen, and  $\chi_j$  ( $\mu\text{mol e}^- \text{ mmol}^{-1} \text{ N s}^{-1}$ ) is the electron transport capacity per unit electron transport nitrogen.  $\chi_{Cj}$  ( $\text{mmol Chl mmol}^{-1} \text{ N}$ ) and  $\chi_C$  ( $\text{mmol Chl mmol}^{-1} \text{ N}$ ) are the conversion coefficients for chlorophyll per electron transport nitrogen and per light harvesting component nitrogen, respectively. Photosynthetic nitrogen partitioning fraction of a pool  $X$  ( $p_X$ ) is determined as the ratio of nitrogen in the pool  $X$  ( $N_X$ ,  $\text{mmol N m}^{-2}$ ) to  $N_{\text{ph}}$ :

$$p_X = N_X / N_{\text{ph}} \quad (3)$$

The rate of change of  $N_X$  is determined by the instantaneous protein synthesis rate [ $S_X(t)$ ,  $\text{mmol N m}^{-2} \text{ }^\circ\text{Cd}^{-1}$ ] and degradation rate [ $D_X(t)$ ,  $\text{mmol N m}^{-2} \text{ }^\circ\text{Cd}^{-1}$ ] of the corresponding enzymes and protein complexes at a given leaf age ( $t$ ,  $^\circ\text{Cd}$ ):

$$dN_X / dt = S_X(t) - D_X(t) \quad (4)$$

Protein synthesis as an age-dependent and zero-order process (Li *et al.*, 2017) is described by a logistic function and independent of the current  $N_X$  state:

$$S_X(t) = 2S_{\text{max},X} / [1 + \exp(t \times t_{d,X})] \quad (5)$$

where  $S_{\text{max},X}$  ( $\text{mmol N m}^{-2} \text{ }^\circ\text{Cd}^{-1}$ ) is the maximum protein synthesis rate of  $N_X$  that occurs at the early stage of leaf development (Supplementary Fig. S1 at JXB online). The constant  $t_{d,X}$  ( $^\circ\text{Cd}^{-1}$ ) describes the relative decreasing rate of the protein synthesis over time (see Table 1 for the coefficients used in the protein turnover model). At age of  $1/t_{d,X}$ ,  $S_X$  reduces to 53.8% of  $S_{\text{max},X}$ .

The degradation rate  $D_X$  is governed by first-order kinetics (Verkroost and Wassen, 2005; Li *et al.*, 2017) with a degradation constant  $D_{r,X}$  ( $^\circ\text{Cd}^{-1}$ ):

$$D_X(t) = D_{r,X} \times N_X(t) \quad (6)$$

The variable  $S_{\text{max},X}$  in Eq. (5) is a function of daily light interception ( $I_{\text{Ld}}$ ,  $\text{mol photons m}^{-2} \text{ d}^{-1}$ ):

$$S_{\text{max},X} = [S_{\text{mm},X} \times k_{i,X} \times I_{\text{Ld}} / (S_{\text{mm},X} + k_{i,X} \times I_{\text{Ld}})] \times r_{N,X} \quad (7)$$

where  $S_{\text{mm},X}$  ( $\text{mmol N m}^{-2} \text{ }^\circ\text{Cd}^{-1}$ ) is the potential maximum protein synthesis rate and  $k_{i,X}$  is the rate constant describing the increase of  $S_{\text{max},X}$  with  $I_{\text{Ld}}$ . The factor  $r_{N,X}$  increases with nitrogen level in the nutrient solution ( $N_S$ , mM) by a Michaelis–Menten constant,  $k_{N,X}$  (mM):

$$r_{N,X} = N_S / (k_{N,X} + N_S) \quad (8)$$

### Modelling leaf photosynthesis

Photosynthetic parameters  $V_{\text{cmax}}$ ,  $J_{\text{max}}$ , and  $Chl$  were estimated from functional nitrogen pools  $N_V$ ,  $N_J$ , and  $N_C$ , using Eq. (2a–c). The net photosynthetic rate ( $A$ ,  $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ ) is defined as the minimum of RuBP carboxylation-limited ( $A_c$ ,  $\text{mmol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ ) and RuBP regeneration-limited ( $A_j$ ,  $\text{mmol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ ) net photosynthetic rate (Farquhar *et al.*, 1980):

$$A = \min(A_c, A_j) \quad (9a)$$

$$A_c = V_c \times (C_c - \Gamma^*) / [C_c + K_c (1 + O / K_o)] - R_d \quad (9b)$$

$$A_j = J \times (C_c - \Gamma^*) / (4C_c + 8\Gamma^*) - R_d \quad (9c)$$

where  $C_c$  ( $\mu\text{mol CO}_2 \text{ mol}^{-1}$ ) is the chloroplastic  $\text{CO}_2$  concentration,  $\Gamma^*$  ( $\mu\text{mol CO}_2 \text{ mol}^{-1}$ ) is the  $\text{CO}_2$  compensation point in the absence of dark respiration,  $K_c$  ( $\mu\text{mol CO}_2 \text{ mol}^{-1}$ ) and  $K_o$  ( $\text{mmol O}_2 \text{ mol}^{-1}$ ) are Michaelis–Menten constants of Rubisco for  $\text{CO}_2$  and  $\text{O}_2$ , respectively,  $O$  ( $\text{mmol O}_2 \text{ mol}^{-1}$ ) is the  $\text{O}_2$  concentration at the site of carboxylation,  $V_c$  ( $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ ) is carboxylation rate, and  $J$  ( $\mu\text{mol e}^- \text{ m}^{-2} \text{ s}^{-1}$ ) is electron transport rate. Daytime respiration rate  $R_d$  ( $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ ) is assumed to vary with  $t$  and the mean  $I_{\text{Ld}}$  during the previous 4 d ( $I_{\text{Ld4d}}$ ):

$$R_d(t) = R_{\text{max}} \times I_{\text{Ld4d}} \times \exp(-R_g \times I_{\text{Ld4d}} \times t) + R_m \times I_{\text{Ld4d}} \times t \quad (10)$$

where  $R_{\text{max}}$  ( $\mu\text{mol CO}_2 \text{ d mol}^{-1} \text{ photons s}^{-1}$ ) relates  $I_{\text{Ld4d}}$  to the maximum  $R_d$ ,  $R_g$  ( $\text{m}^2 \text{ d }^\circ\text{Cd}^{-1} \text{ mol}^{-1} \text{ photons}$ ) influences the decrease in the growth respiration, and  $R_m$  ( $\mu\text{mol CO}_2 \text{ d }^\circ\text{Cd}^{-1} \text{ mol}^{-1} \text{ photons s}^{-1}$ ) affects the increase in the maintenance respiration with  $t$ .

$V_c$  and  $J$  are calculated from  $V_{\text{cmax}}$  and  $J_{\text{max}}$ , respectively, depending on the photosynthetic photon flux density (PPFD) incident on the leaf ( $I_{\text{Lc}}$ ,  $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ ) according to Qian *et al.* (2012) and Ögren and Evans (1993), respectively:

$$V_c = V_{\text{cmax}} \left\{ 0.31 + \frac{0.69}{1 + \exp[-0.009 \times (I_{\text{Lc}} - 500)]} \right\} \quad (11)$$

$$J = \left\{ \phi \times \alpha \times I_{\text{Lc}} + J_{\text{max}} - \left[ \frac{(\phi \times \alpha \times I_{\text{Lc}} + J_{\text{max}})^2}{-4\theta \times J_{\text{max}} \times \phi \times \alpha \times I_{\text{Lc}}} \right]^{0.5} \right\} / (2\theta) \quad (12)$$

where  $\phi$  ( $\mu\text{mol e}^- \mu\text{mol photons}^{-1}$ ) is the conversion efficiency of photons to  $J$ , and  $\theta$  (unitless) is a constant convexity factor describing the response of  $J$  to  $I_{\text{Lc}}$ . Leaf absorptance ( $\alpha$ , unitless) is related to  $Chl$  (Evans, 1993):

$$\alpha = Chl / (Chl + 0.076) \quad (13)$$

**Table 1.** List of coefficients used in the protein turnover model for photosynthetic nitrogen pools, carboxylation pool  $N_V$ , electron transport pool  $N_J$ , and light harvesting pool  $N_C$

Description	Coefficient	Unit	Pool $N_V$	Pool $N_J$	Pool $N_C$
Degradation constant [Eq. (6)]	$D_r$	$^\circ\text{Cd}^{-1}$	0.0195	0.0195	0.0091
Increase rate constant of $S_{\text{max}}$ with $I_{\text{Ld}}$ [Eq. (7)]	$k_i$	$\text{mmol N m}^2 \text{ ground d m}^{-2}$ LA $^\circ\text{Cd}^{-1} \text{ mol}^{-1} \text{ photon}$	0.173	0.130	0.234
Michaelis–Menten constant relating $N_S$ to $S_{\text{max}}$ [Eq. (8)]	$k_N$	mM	0.536	0.420	0.316
Potential maximum synthesis rate [Eq. (7)]	$S_{\text{mm}}$	$\text{mmol N m}^{-2} \text{ }^\circ\text{Cd}^{-1}$	1.122	0.852	0.248
Decreasing constant of synthesis rate [Eq. (5)]	$t_d$	$^\circ\text{Cd}^{-1}$	0.001	0.002	0.001

The coefficients were estimated from the growth chamber experiment. Model variables and other coefficients are listed in Tables 2 and 3.

**Table 2.** List of model input and output variables

Description	Variable	Unit	Equation	Type
Net photosynthetic rate	$A$	$\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$	9a	Output
RuBP carboxylation-limited $A$	$A_c$	$\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$	9b	Output
RuBP regeneration-limited $A$	$A_j$	$\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$	9c	Output
Leaf absorptance	$\alpha$	—	13	Output
Atmospheric $\text{CO}_2$ concentration	$C_a$	$\mu\text{mol CO}_2 \text{ mol}^{-1}$	—	Input
Chloroplastic $\text{CO}_2$ concentration	$C_c$	$\mu\text{mol CO}_2 \text{ mol}^{-1}$	14	Output
Leaf chlorophyll per unit area	$Chl$	$\text{mmol m}^{-2}$	2c	Output
Leaf-to-air vapor pressure deficit	$D$	kPa	—	Input
Protein degradation rate of N pool $X$	$D_X$	$^{\circ}\text{Cd}^{-1}$	6	Output
Factor for creating variation in N distribution	$f_d$	—	18	Input
Factor for creating variation in N partitioning	$f_p$	—	19	Input
Mesophyll conductance to $\text{CO}_2$	$g_m$	$\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$	16	Output
Maximum $g_m$	$g_{m\text{max}}$	$\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$	17	Output
Stomatal conductance to $\text{CO}_2$	$g_{sc}$	$\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$	15	Output
PPFD at leaf	$I_{Lc}$	$\mu\text{mol photons m}^{-2} \text{ s}^{-1}$	—	Input
Daily photosynthetic photon integral at leaf	$I_{Ld}$	$\text{mol photons m}^{-2} \text{ d}^{-1}$	—	Input
Mean $I_{Ld}$ during the last 4 d	$I_{Ld4d}$	$\text{mol photons m}^{-2} \text{ d}^{-1}$	—	Input
Electron transport rate	$J$	$\mu\text{mol e}^- \text{ m}^{-2} \text{ s}^{-1}$	12	Output
Maximum electron transport rate	$J_{\text{max}}$	$\mu\text{mol e}^- \text{ m}^{-2} \text{ s}^{-1}$	2b	Output
Leaf area	LA	$\text{m}^2$	—	Input
Total leaf photosynthetic N content in the canopy	$N_{\text{canopy}}$	mmol N	—	Output
Leaf photosynthetic N content	$N_{\text{leaf}}$	mmol N	—	Output
Leaf photosynthetic N per unit area	$N_{\text{ph}}$	mmol N $\text{m}^{-2}$	1	Output
N concentration of nutrient solution	$N_s$	mM	—	Input
Concentration of N pool $X$	$N_X$	mmol N $\text{m}^{-2}$	4	Output
Concentration of N pool of light harvesting	$N_C$	mmol N $\text{m}^{-2}$	4	Output
Concentration of N pool of electron transport	$N_J$	mmol N $\text{m}^{-2}$	4	Output
Concentration of N pool of carboxylation	$N_V$	mmol N $\text{m}^{-2}$	4	Output
Partitioning fraction of N pool $X$	$p_X$	—	3	Output
Daytime respiration rate in the absence of photorespiration	$R_d$	$\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$	10	Output
Reduction factor of protein synthesis (depending on N availability)	$r_N$	—	8	Output
Maximum protein synthesis rate	$S_{\text{max}}$	mmol N $\text{m}^{-2} \text{ }^{\circ}\text{Cd}^{-1}$	7	Output
Protein synthesis rate of N pool $X$	$S_X$	mmol N $\text{m}^{-2} \text{ }^{\circ}\text{Cd}^{-1}$	5	Output
Leaf age	$t$	$^{\circ}\text{Cd}$	—	Input
Carboxylation rate	$V_c$	$\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$	11	Output
Maximum carboxylation rate	$V_{c\text{max}}$	$\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$	2a	Output

Chloroplastic  $\text{CO}_2$  concentration depends on the steady-state of stomatal conductance ( $g_{sc}$ ,  $\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ ) and mesophyll conductance ( $g_m$ ,  $\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ ) to  $\text{CO}_2$ :

$$C_c = C_a - A \times \left[ \frac{(g_{sc} + g_m)}{(g_{sc} \times g_m)} \right] \quad (14)$$

where  $C_a$  ( $\mu\text{mol CO}_2 \text{ mol}^{-1}$ ) is atmospheric  $\text{CO}_2$  concentration, and  $g_{sc}$  is calculated with species-specific constants of stomatal conductance,  $g_0$  and  $g_1$  (Chen *et al.*, 2014), and leaf-to-air vapor pressure deficit ( $D$ , kPa, Medlyn *et al.*, 2011):

$$g_{sc} = g_0 + \left(1 + g_1 / \sqrt{D}\right) \times A / C_a \quad (15)$$

Mesophyll conductance is expressed as a log-normal function of  $t$  (Chen *et al.*, 2014), where  $g_m$  first increases during leaf development and decreases during ageing (Flexas *et al.*, 2008):

$$g_m = g_{m\text{max}} \times \exp \left\{ -0.5 \times \left[ \ln \left( t / t_{g_m} \right) / v_{g_m} \right]^2 \right\} \quad (16)$$

where  $t_{g_m}$  is the  $t$  when the maximum  $g_m$  ( $g_{m\text{max}}$ ,  $\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ ) occurs and  $v_{g_m}$  is the standard deviation of the curve;  $g_{m\text{max}}$  is linearly related to  $N_{\text{ph}}$ , since a similar relationship has been reported for  $C_3$  plants (e.g. Yamori *et al.*, 2011a):

$$g_{m\text{max}} = r_{g_m} \times N_{\text{ph}} + r_{g_m0} \quad (17)$$

where  $r_{g_m}$  ( $\text{mol CO}_2 \text{ mmol}^{-1} \text{ N s}^{-1}$ ) describes the rate of increase of  $g_{m\text{max}}$  in relation to  $N_{\text{ph}}$ , and  $r_{g_m0}$  ( $\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ ) is the minimum  $g_{m\text{max}}$ .

The steady-state  $A_c$  was solved analytically with Eqs (9b), (14), and (15), and  $A_j$  with Eqs (9c), (14), and (15), following Moualeu-Ngangue *et al.* (2016). Model variables and coefficients are listed in Tables 1–3.

#### Growth chamber experiment to investigate the dynamics of photosynthetic protein turnover

Cucumber (*Cucumis sativus* ‘Aramon’, Rijk Zwaan, De Lier, The Netherlands) plants were grown in two experiments at the Institute of Horticultural Production Systems, Leibniz Universität Hannover, Germany (latitude 52.4°N).

One growth chamber experiment was conducted from 21 October to 9 December 2016 with factorial combinations of three light and three nitrogen supply levels to parameterize the photosynthetic protein turnover model (see below). Cucumber seeds were sown in rock-wool cubes (36 × 36 × 40 mm) on 5 October. Eight days later, seedlings were transplanted to larger rock-wool cubes (10 × 10 × 6.2 cm) for another 8 d until the second true leaves appeared (leaf length ≥ 3 cm). Plants were transferred into 25 litre plastic containers (one plant per container) on 21 October and cultivated hydroponically with a 12 h light period and

**Table 3.** List of model coefficients

Description	Coefficient	Unit	Value (SE)	Reference
Conversion coefficient of chlorophyll per light harvesting N	$\chi_C$	mmol Chl mmol <sup>-1</sup> N	0.03384	Buckley <i>et al.</i> (2013)
Conversion coefficient of chlorophyll per electron transport N	$\chi_{CJ}$	mmol Chl mmol <sup>-1</sup> N	$4.64 \times 10^{-4}$	Buckley <i>et al.</i> (2013)
Conversion coefficient of electron transport capacity per electron transport N	$\chi_J$	$\mu\text{mol e}^- \text{mmol}^{-1} \text{N s}^{-1}$	9.48	Buckley <i>et al.</i> (2013)
Conversion coefficient of carboxylation capacity per Rubisco N	$\chi_V$	$\mu\text{mol CO}_2 \text{mmol}^{-1} \text{N s}^{-1}$	4.49	Buckley <i>et al.</i> (2013)
Minimum $g_{sc}$	$g_0$	$\text{mol CO}_2 \text{m}^{-2} \text{s}^{-1}$	0.009	Chen <i>et al.</i> (2014)
Species-specific coefficient of $g_{sc}$	$g_1$	—	3.51	Chen <i>et al.</i> (2014)
CO <sub>2</sub> compensation point in the absence of dark respiration	$\Gamma^*$	$\mu\text{mol CO}_2 \text{mol}^{-1}$	43.02	Singsaas <i>et al.</i> (2004)
Michaelis–Menten constant of Rubisco for CO <sub>2</sub>	$K_C$	$\mu\text{mol CO}_2 \text{mol}^{-1}$	404	Chen <i>et al.</i> (2014)
Michaelis–Menten constant of Rubisco for O <sub>2</sub>	$K_O$	mmol O <sub>2</sub> mol <sup>-1</sup>	278	Chen <i>et al.</i> (2014)
O <sub>2</sub> concentration at the site of carboxylation	$O$	mmol O <sub>2</sub> mol <sup>-1</sup>	210	Chen <i>et al.</i> (2014)
Coefficient relating $N_{ph}$ to $g_{mmax}$	$r_{gm}$	$\text{mol CO}_2 \text{mmol}^{-1} \text{N s}^{-1}$	$1.64 \times 10^{-3}$ ( $5.27 \times 10^{-4}$ )	—
Minimum $g_{mmax}$	$r_{gm0}$	$\text{mol CO}_2 \text{m}^{-2} \text{s}^{-1}$	0.140 (0.0345)	—
Coefficient related to the decrease in $R_d$ by growth respiration	$R_g$	$\text{m}^2 \text{d} \text{ } ^\circ\text{Cd}^{-1} \text{mol}^{-1} \text{photon}$	$4.16 \times 10^{-4}$ ( $4.52 \times 10^{-5}$ )	—
Coefficient related to the increase in $R_d$ by maintenance respiration	$R_m$	$\mu\text{mol CO}_2 \text{d} \text{ } ^\circ\text{Cd}^{-1} \text{mol}^{-1} \text{photons s}^{-1}$	$1.88 \times 10^{-4}$ ( $1.61 \times 10^{-5}$ )	—
Coefficient relating $I_{Ld}$ to maximum $R_d$	$R_{max}$	$\mu\text{mol CO}_2 \mu\text{mol}^{-1} \text{photons s}^{-1}$	0.308 (0.028)	—
Conversion efficiency of photons to $J$	$\phi$	$\mu\text{mol e}^- \mu\text{mol}^{-1} \text{photons}$	$0.340 (2.5 \times 10^{-3})$	—
Convexity coefficient	$\theta$	—	0.7	Chen <i>et al.</i> (2014)
Leaf age when $g_{mmax}$ occurs	$t_{gm}$	$^\circ\text{Cd}$	121 (8.1)	—
Standard deviation of the dependence of $g_m$ - $t$ curve	$v_{gm}$	—	0.860 (0.063)	—

Standard errors (SE) are indicated in parentheses.

24 °C day/20 °C night air temperature. Three nitrogen levels, 9.6, 4.6 and 2.3 mM, were supplied using Ca(NO<sub>3</sub>)<sub>2</sub> and Fertyl Basisdünger 1 (Planta GmbH, Regenstauf, Germany, 5.2 mM K, 1.3 mM P, 0.82 mM Mg in working solution). Nutrient solution was replaced weekly and adjusted to pH 6.0–6.5 two times a week. Three constant light conditions with daily photosynthetic photon integrals (DPI) of 28.9, 14.2, and 4.4 mol photons m<sup>-2</sup> d<sup>-1</sup> were provided using metal halide lamps. Four plants were grown under each treatment combination. Three leaves per plant (between leaf ranks four to eight, counted acropetally) were maintained horizontally and well exposed to incoming light using custom-made leaf holders, while the rest of the shoot was trained downward to avoid mutual shading. Gas exchange (see below) and relative chlorophyll content (SPAD-502; Minolta Camera, Japan) were measured at different thermal ages of the leaves, ranging from 45 °Cd to 558 °Cd, calculated by subtracting a base temperature of 10 °C (Savvides *et al.*, 2016) from mean daily air temperature around the leaf. Air temperature was recorded continuously using data loggers (Tinytag; Gemini Data Loggers, Chichester, UK). After gas exchange measurements, leaves were harvested for leaf area and nitrogen analyses.

#### Greenhouse experiment to evaluate optimality of nitrogen distribution and partitioning

One greenhouse experiment was carried out from 4 April to 12 May 2017 under two light regimes and two nitrogen supply levels to evaluate the model performance and to collect input data for optimality analyses. Seeds were sown on 14 March and transplanted to larger rock-wool cubes on 22 March. After the third true leaves had appeared, plants were transferred onto rock-wool slabs on 4 April with plant density of 1.33 plants m<sup>-2</sup> and supplied with two nitrogen concentrations, 10 mM (high nitrogen, HN) and 2.5 mM (low nitrogen, LN), by drip irrigation using the same fertilizers as described in the growth chamber experiment. During the experimental period, average nitrogen supply was calculated from the nitrogen concentration in the nutrient supply and rock-wool slabs, which was 8.2 and 2.0 mM for HN and LN, respectively. Plants were grown

under either high light (HL) or low light (LL) regimes. The southern half of the greenhouse was unshaded as the HL regime. The LL regime was created in the northern half of the greenhouse by shading nets to reduce incoming light from top and sides, where PPFD was reduced on average to ca. 40% of that under HL ( $38 \pm 1.3\%$  under sunny and  $42 \pm 0.2\%$  under cloudy condition). Average DPI above the canopy was 21.4 and 8.5 mol photons m<sup>-2</sup> d<sup>-1</sup> for HL and LL, respectively, during the experimental period. DPI during the experimental period was recorded by the weather station located above the greenhouse. An average light transmittance of 49.8% through the greenhouse structure was applied (39.2% on a sunny day and 60.4% on a cloudy day). Air temperature in the middle canopy was recorded continuously using data loggers and was significantly higher under HL (0.5 °Cd per day). Gas exchange measurements and harvests were conducted at four time points on 21 April, 28 April, 5 May, and 12 May at two different canopy layers with two replications. Leaf age at measurement ranged from 77 to 414 °Cd. Leaf elevation angle was obtained by a 3D digitizer (Fastrak; Polhemus, Colchester, VT, USA) according to Chen *et al.* (2014). Leaves were harvested after gas exchange measurements to determine leaf area index (LAI, m<sup>2</sup> m<sup>-2</sup>).

#### Gas exchange measurements and estimation of photosynthetic parameters

Light-saturated net photosynthetic rate under PPFD of 1300  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  ( $A_{1300}$ ,  $\mu\text{mol CO}_2 \text{m}^{-2} \text{s}^{-1}$ ) and light response curves were measured using a portable photosynthesis system (LI-6400XT; Li-Cor Inc., Lincoln, NE, USA). All measurements were carried out under sample CO<sub>2</sub> 400  $\mu\text{mol mol}^{-1}$ , leaf temperature 25 °C and relative humidity 55–65%.  $R_d$  was estimated from the linear portion of the light response curve (Kok, 1948).  $V_{cmax}$  was estimated using the one-point method (Wilson *et al.*, 2000; De Kauwe *et al.*, 2016), and  $J_{max}$  and  $\phi$  by least squares fitting to a non-rectangular hyperbola (Ögren and Evans, 1993). Mesophyll conductance was estimated using the variable  $J$  method (Harley *et al.*, 1992). Chlorophyll fluorescence was measured using the multiphase flash approach (Loriaux *et al.*, 2013) following Moualeu-Ngangue *et al.* (2017).

### Nitrogen analyses and photosynthetic nitrogen partitioning estimation

Leaf samples obtained in the growth chamber experiment were freeze-dried and ground into a fine powder for nitrogen analyses. Total leaf nitrogen was analysed using the Kjeldahl method (Nelson and Sommers, 1980). Leaf chlorophyll was extracted with 96% ethanol and analysed colorimetrically (Lichtenthaler, 1987). Relationships between relative chlorophyll content (SPAD) and *Chl* were determined (Supplementary Fig. S2) for estimating *Chl* in the greenhouse experiment.

### Model parameterization

The differential equations (4)–(6) were solved and the coefficients were quantified using R (version 3.3.0; R Foundation for Statistical Computing, Vienna, Austria) by using the packages ‘deSolve’ and ‘DEoptim’, which minimizes the sums of squares of the residuals between observations and simulations. The data obtained in the growth chamber experiment were used for the parameterization.  $D_{r,X}$  and  $t_{d,X}$  were first quantified for each pool using data of all treatments. With the determined values of  $D_{r,X}$  and  $t_{d,X}$ ,  $S_{\max,X}$  was then quantified for each treatment.  $S_{\max,X}$ ,  $k_{I,X}$ , and  $k_{N,X}$  were determined from  $S_{\max,X}$  [Eqs (7) and (8)] by least squares fitting in SigmaPlot (version 11.0, Systat software GmbH, Erkrath, Germany) as well as the influences of  $t$  and  $I_{Ld}$  on  $R_d$  [Eq. (10)] and  $g_m$  [Eqs (16) and (17)].

### Dynamic leaf photosynthetic nitrogen simulation and model evaluation

Daily environmental information during the experimental period (Supplementary Fig. S3) and the canopy information obtained at the four harvests, including age and area of each leaf, were used as input to simulate photosynthetic nitrogen per unit leaf area ( $N_{ph}$ , mmol N m<sup>-2</sup>), photosynthetic nitrogen per leaf ( $N_{leaf}$ , mmol N) and total leaf photosynthetic nitrogen content of the canopy ( $N_{canopy}$ , mmol N). First, leaf elevation angle of each leaf and LAI were simulated empirically depending on  $t$  (Supplementary Fig. S4). Second, for each time step, the daily light interception  $I_{Ld}$  at the leaf was calculated and used in Eq. (7) to simulate protein turnover. Light interception was calculated by the Beer–Lambert law (Monsi and Saeki, 1953) with a light extinction coefficient of 0.695 and adjusted by the cosine of leaf elevation angle. For model evaluation, root mean squared deviation (RMSD) and accuracy (%) were determined for photosynthetic parameters,  $N_{ph}$ , and  $p_X$  predictions following Kahlen and Stützel (2011).

### Simulating daily canopy carbon assimilation

Daily canopy carbon assimilation during daytime (DCA, mol CO<sub>2</sub> d<sup>-1</sup>) was simulated using greenhouse canopy characteristics obtained at the last harvest as input (Supplementary Table S1; Supplementary Fig. S5). Leaf-to-air vapor pressure deficit ( $D$ ) 1.2 kPa and  $C_a$  400 μmol CO<sub>2</sub> mol<sup>-1</sup> were used in all simulations, similar to the environmental conditions during the gas exchange measurements. Scenarios with different DPI levels were defined for simulating DCA. Up to six DPI levels were taken as relative to the average DPI during acclimation (aDPI) to simulate the influence of day-to-day DPI fluctuation on DCA. To simulate DCA, diurnal PPFD above the canopy was simulated for a given DPI level with a time step of 0.1 h by a simple cosine bell function (Kimball and Bellamy, 1986) with 14.4 h day length.

### Modifying photosynthetic nitrogen distribution and partitioning

To evaluate the effects of between-leaf distribution and within-leaf partitioning of  $N_{ph}$  on DCA, a distribution factor  $f_d$  was introduced into Eq. (5) to create variations in the rate of protein synthesis, and a partitioning factor  $f_{p,X}$  was introduced into Eq. (7) to create variations in the maximum protein synthesis rate of different functional pools:

$$S_X(t) = 2S_{\max,X} / \left[ 1 + \exp(t \times t_{d,X} \times f_d) \right] \quad (18)$$

$$S_{\max,X} = \left[ \frac{S_{\max,X} \times f_{p,X} \times k_{I,X} \times I_{Ld}}{(S_{\max,X} \times f_{p,X} + k_{I,X} \times I_{Ld})} \right] \times t_{N,X} \quad (19)$$

A control condition was defined with  $f_d=1$  and  $f_{p,X}=1$ , when all coefficients in the synthesis process (Table 1) remained unmodified. Increasing  $f_d$  accelerates the decrease in the rate of protein synthesis and enhances acropetal  $N_{ph}$  reallocation. An increase in  $f_{p,X}$  results in a higher rate of synthesis of  $N_X$  and increases the partitioning to pool  $X$ . A modified partitioning pattern that maximized DCA was identified as optimal for several DPI levels, and the optimal values of  $f_{p,X}$  were determined using the package ‘DEoptim’ in R. The change in DCA caused by modified distribution or optimal partitioning of  $N_{ph}$  was compared with the control conditions. The ratios between optimal and control partitioning fractions of each pool  $X$ , as well as the contributions of daily leaf carbon assimilation (DLA) to the DCA increase were calculated along the canopy depth.


## Results

### Mechanistic model aims to quantify the environmental effects of light and nitrogen availabilities and developmental effects on photosynthetic protein turnover

In the model, we assume that photosynthetic protein turnover is under genetic and environmental control. The genetic control is characterized by the potential maximum protein synthesis rate  $S_{\max}$ , coefficient  $t_d$ , and protein degradation constant,  $D_r$ . The coefficient  $t_d$  affects the decrease in the rate of synthesis, and  $D_r$  contributes to the degradation rate, which together influence the developmental effect on protein turnover dynamics. The low value of  $t_d$  (0.001–0.002 °Cd<sup>-1</sup>, Table 1) suggests that the influence of ageing appears rather late in the leaf lifespan under a constant light environment. The coefficient  $D_r$  was found to be the same for the carboxylation pool ( $N_V$ ) and the electron transport pool ( $N_J$ ), while the light harvesting pool ( $N_C$ ) had a lower  $D_r$  (Table 1). The genotypic sensitivities to light and nitrogen availabilities are characterized by  $k_I$  and  $k_N$ , respectively. Collectively,  $S_{\max}$ ,  $k_I$ , and  $k_N$  determine the maximum protein synthesis rate  $S_{\max}$  in Eq. (7). When light was increased 2.5-fold (from LL to HL),  $S_{\max}$  increased by 50% in  $N_V$  and  $N_J$ , and by 10% in  $N_C$ , while nitrogen level had less influence on  $S_{\max}$  (<10%), which only occurred under low nitrogen concentration (<3.5 mM) and the higher light intensity (Fig. 1), showing that light had a major control of  $S_{\max}$ .  $N_C$  had the highest  $k_I$  (Table 1); consequently,  $S_{\max,C}$  approached saturation at lower light intensity than  $S_{\max,V}$  and  $S_{\max,J}$  (Fig. 1).  $S_{\max,V}$  and  $S_{\max,J}$  were well coordinated in response to light and nitrogen level (Fig. 1A, B), but the higher  $k_I$  and  $k_N$  of  $N_V$  (Table 1) suggested that  $N_V$  synthesis is more sensitive to the variation in light and nitrogen availabilities than  $N_J$ .

### Effects of light and nitrogen availabilities on maximal protein synthesis rate explain the dynamics of photosynthetic acclimation

We evaluated the model using a greenhouse experiment, where leaves grown under combinations of two light regimes (HL and LL) and two nitrogen levels (HN and LN) were measured



**Fig. 1.** Simulated effects of daily light interception ( $I_{Ld}$ , mol photons  $m^{-2} d^{-1}$ ) and nitrogen supply level in the nutrient solution ( $N_S$ , mM) on maximum protein synthesis rate ( $S_{max,X}$ ) in Eq. (7) using coefficients from Table 1, of (A) the carboxylation, (B) the electron transport and (C) the light harvesting pools. The colors denote the normalized maximum protein synthesis rate, which is  $S_{max,X}$  normalized by the potential maximum protein synthesis rate ( $S_{mm,X}$ ) in Eq. (7). The data obtained in the growth chamber experiment were used for the parameterization. The arrows above and beside the figures indicate the corresponding average environmental conditions in the greenhouse experiment: high light (HL) 21.4 mol photons  $m^{-2} d^{-1}$ ; low light (LL) 8.5 mol photons  $m^{-2} d^{-1}$ ; high nitrogen (HN) 8.2 mM; low nitrogen (LN) 2.0 mM.

in two canopy layers weekly for four consecutive weeks. The model predicted leaf photosynthetic characteristics with high accuracy (70–91%, Fig. 2) and a trend of photosynthetic acclimation (Supplementary Fig. S6) similar to the experimental observations (Fig. 3), except for slight overestimations in photosynthetic nitrogen ( $N_{ph}$ , Fig. 2C), carboxylation pool (Fig. 2D, G), and chlorophyll (Fig. 2F).

Photosynthetic acclimation in the greenhouse canopies as influenced by the interplay between light, nitrogen level and leaf age was examined (Fig. 3). Light had positive effects on  $N_{ph}$  (Fig. 3A), the partitioning fractions of  $N_V$  ( $p_V$ , Fig. 3C) and  $N_J$  ( $p_J$ , Fig. 3E) but negative effects on the partitioning fraction of  $N_C$  ( $p_C$ , Fig. 3G). This negative effect of light on  $p_C$  can be explained by the high  $k_I$  of  $N_C$  (Table 1), which leads to an saturation of  $S_{max,C}$  under lower light (Fig. 1). The changes in  $N_{ph}$ ,  $p_V$ ,  $p_J$ , and  $p_C$  with leaf age were similar to those with light (Fig. 3) due to the association in the gradients of age and light.

In comparison with HN,  $N_{ph}$  under LN was significantly lower in the young leaves but similar in the old leaves (Fig. 3A). In the greenhouse, young leaves developed under high light intensity, which increased the sensitivity of  $S_{max}$  to nitrogen level (Fig. 1). During the simultaneous increase in leaf age and mutual shading, the effects of nitrogen supply on  $S_{max}$  became less prevalent (Fig. 1). Nitrogen level had less influence on functional partitioning (Fig. 3C, E, G) than light (Fig. 3D, F, H).

*Photosynthetic nitrogen distribution is close to optimum and the effect of nitrogen reallocation is more prominent under limited nitrogen availability*

The influence of  $N_{ph}$  distribution pattern along the canopy depth on daily canopy carbon assimilation (DCA, mol  $CO_2 d^{-1}$ ) was evaluated by introducing a distribution factor  $f_d$  to create variations in the rate of protein synthesis. In our model, protein synthesis and degradation rates determined simultaneously (i) total leaf photosynthetic nitrogen content of the canopy ( $N_{canopy}$ , mmol N), (ii)  $N_{ph}$  distribution in the canopy, and (iii)

$N_{ph}$  partitioning fractions of pools  $X$  ( $p_X$ ) in the leaf. Thus, it was impossible to modify single elements while maintaining the other two constant. Increasing  $f_d$  led to a faster reduction of  $N_{ph}$  during leaf ageing and more acropetal  $N_{ph}$  reallocation. However, it also reduced  $N_{canopy}$  and tended to increase  $p_C$  (data not shown). Therefore, to obtain the leaf photosynthetic nitrogen content ( $N_{leaf,i}$ , mmol N in leaf  $i$ ) with comparable  $N_{canopy}$ , simulated  $N_{leaf,i}$  with  $f_d=n$  (denoted as  $N'_{leaf,i}$ ) was adjusted proportionally to the ratio between  $N_{canopy}$  calculated with  $f_d=1$  and with  $f_d=n$ :


$$N_{leaf,i}(f_d = n) = N'_{leaf,i}(f_d = n) \times \left[ N_{canopy}(f_d = 1) / N_{canopy}(f_d = n) \right] \quad (20a)$$

$p_{X,i}$  was set equal to the control value:

$$p_{X,i} = N_{X,i}(f_d = 1) / N_{ph,i}(f_d = 1) \quad (20b)$$

These adjustments assured the same amount of  $N_{canopy}$  among the distribution patterns. The factor  $f_d$  was varied between 0.5 and 5.0 at intervals of 0.5 in the simulation, which gave values of  $N_{ph}$  comparable to those measured in cucumber leaves (22–135 mmol N  $m^{-2}$ ; Fig. 4). Canopy  $N_{ph}$  distributions with enhanced acropetal reallocation were created by increasing  $f_d$  (Fig. 4; Supplementary Fig. S7). In general, the distribution of  $N_{ph}$  corresponded to the vertical light distribution except in the expanding leaves in the upper canopy, and the  $N_{ph}$  distribution with light was steeper under LL (Supplementary Fig. S7).

To simulate the natural fluctuations in light between days, three light levels representing 200% (aDPI<sub>200</sub>), 100% (aDPI<sub>100</sub>) and 50% (aDPI<sub>50</sub>) of average DPI during acclimation (aDPI) were used in the DCA simulation. Under aDPI<sub>100</sub> and aDPI<sub>50</sub>, enhancing acropetal  $N_{ph}$  reallocation did not significantly increase DCA (<5%), suggesting that  $N_{ph}$  distribution was optimal under constant and decreasing DPI (Fig. 5B, C). More acropetal reallocation did not improve the optimality in  $N_{ph}$  distribution in terms of maximizing DCA since a large proportion of leaf area was located in the middle-lower to lower



**Fig. 2.** Comparisons between simulated and observed leaf photosynthetic parameters. (A) Light-saturated net photosynthetic rate under PPFD 1300  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  ( $A_{1300}$ ,  $\mu\text{mol CO}_2 \text{m}^{-2} \text{s}^{-1}$ ); (B) daytime respiration rate ( $R_d$ ,  $\mu\text{mol CO}_2 \text{m}^{-2} \text{s}^{-1}$ ); (C) leaf photosynthetic nitrogen ( $N_{\text{ph}}$ ,  $\text{mmol N m}^{-2}$ ); (D) maximum carboxylation rate ( $V_{\text{cmax}}$ ,  $\mu\text{mol CO}_2 \text{m}^{-2} \text{s}^{-1}$ ); (E) maximum electron transport rate ( $J_{\text{max}}$ ,  $\mu\text{mol e}^- \text{m}^{-2} \text{s}^{-1}$ ); (F) chlorophyll ( $Chl$ ,  $\text{mmol Chl m}^{-2}$ ); (G) partitioning fraction of the carboxylation pool ( $\rho_V$ ); (H) partitioning fraction of the electron transport pool ( $\rho_J$ ); and (I) partitioning fraction of the light harvesting pool ( $\rho_C$ ). The observed data were obtained in the greenhouse experiment. The dotted grey lines are one-to-one lines. Root mean squared deviation (RMSD) and accuracy of the predictions are shown (see Materials and methods).

canopy (Supplementary Fig. S5). However, enhancing  $N_{\text{ph}}$  reallocation resulted in an increase in DCA by 7% under LN at  $a\text{DPI}_{200}$  (Fig. 5A), indicating that acropetal  $N_{\text{ph}}$  reallocation was more important under LN than HN.


It was observed that  $N_{\text{ph}}$  was more overestimated in the older leaves than in the younger ones (Fig. 2C), which indicated that our model tended to underestimate the acropetal  $N_{\text{ph}}$  reallocation when scaling up from leaf to canopy level. In order to maintain a constant light environment for the measured leaves in the growth chamber experiment, leaves younger than the sampled leaves were trained downward and their light interception, together with their nitrogen demand, was inevitably

reduced; therefore, the model coefficients were obtained from the leaves with limited nitrogen reallocation. However, underestimating acropetal  $N_{\text{ph}}$  reallocation would not affect our result that  $N_{\text{ph}}$  distribution was close to optimum.


#### Suboptimal nitrogen partitioning is due to daily light fluctuation

To find the optimal within-leaf  $N_{\text{ph}}$  partitioning between functions, the potential maximal protein synthesis rate for pool X was modified by a factor  $f_{\text{p},X}$ , ranging from 0.2 to 2.0. Increasing  $f_{\text{p},X}$  resulted in higher protein synthesis rates, but it





**Fig. 3.** Comparisons of leaf photosynthetic nitrogen ( $N_{ph}$ ,  $\text{mmol N m}^{-2}$ ; A, B), partitioning fractions of the carboxylation pool ( $p_V$ ; C, D), the electron transport pool ( $p_J$ ; E, F), and the light harvesting pool ( $p_C$ ; G, H) between high and low nitrogen supply (HN and LN, respectively; A, C, E, G) and between high and low light conditions (HL and LL, respectively; B, D, F, H). Each point represents the measurements in the greenhouse experiment obtained from a comparable canopy layer. The orange open circles indicate leaves grown under HL, the black closed circles indicate LL, the blue open squares indicate HN and the black closed squares indicate LN. The size of the circles increases with leaf age, ranging from 77 °Cd to 414 °Cd. The solid lines show the linear regression  $y = ax + b$ . The  $P$  values of the slope  $a$  are shown. The values of  $a$  are specified with 95% confidence intervals when they are significantly different from 1. The dotted grey lines are one-to-one lines.



**Fig. 4.** Leaf photosynthetic nitrogen ( $N_{ph}$ ,  $\text{mmol N m}^{-2}$ ) distributions along the canopy depth, characterized by leaf area index (LAI,  $\text{m}^2 \text{m}^{-2}$ ). Variations in nitrogen distribution were created using a distribution factor  $f_d$  ranging from 0.5 to 5.0 at intervals of 0.5 in Eq. (18) under different growth conditions. (A) High nitrogen and high light (HN+HL); (B) high nitrogen and low light (HN+LL); (C) low nitrogen and high light (LN+HL); (D) low nitrogen and low light (LN+LL). Simulated control  $N_{ph}$  distributions ( $f_d=1$ ) are indicated by the green lines.

also increased  $N_{\text{canopy}}$  and the proportion of nitrogen distributed in the upper canopy. After simulating nitrogen partitioning with a modified  $f_{p,x}$ ,  $N_{\text{leaf}}$  of each leaf was re-assigned to their control values that were obtained with  $f_{p,x}=1$ . Partitioning patterns with maximal DCA at six DPI levels (25–400% aDPI) were identified as optimal and the maximal DCA was compared with control DCA (Fig. 6). The increase in DCA by optimal partitioning was insignificant (<5%) when DPI was close to aDPI (indicated by the arrows in Fig. 6). This suggested the ability of plants to maximize DCA by optimizing  $N_{ph}$  partitioning to aDPI.  $N_{ph}$  partitioning deviated further from optimum when DPI diverged from aDPI (Fig. 6). Therefore, strong day-to-day light fluctuation induced suboptimality in  $N_{ph}$  partitioning and led to lower PNUE.

By optimizing  $N_{ph}$  partitioning, DCA could be increased by nitrogen reinvestment in the limited functional pools. Under aDPI<sub>200</sub>,  $N_{ph}$  partitioning was suboptimal under HL (Fig. 6), and this suboptimality was less under HN than under LN (Table 4). By reinvesting about half of  $N_C$  into  $N_V$  and  $N_J$  (Fig. 7A, C), DCA increased by 6% under HN and by 13% under LN (Table 4), as a result of increased carbon assimilation in the middle-lower canopy (Fig. 7A, C). Under aDPI<sub>50</sub>, HN did not reduce the suboptimality in  $N_{ph}$  partitioning (Table 4) due to an underinvestment in the light harvesting function.


Reinvesting  $N_V$  into  $N_C$  in the middle or upper canopy (HL, Fig. 8A; LL, 8B, 8D) increased DCA by 7–25% (Table 4).

## Discussion

This model is the first approach applying a dynamic protein turnover mechanism at the leaf level to assess the optimality and limitation in nitrogen use at the canopy level. Here, maximized canopy carbon assimilation is considered as a general indicator of maximizing fitness. The adaptation of the protein turnover mechanism gives reasonable predictions of optimal  $N_{ph}$  and accurate predictions of leaf photosynthetic traits.


### *Mechanistic explanation of leaf nitrogen economics under a wide range of light and nitrogen availabilities*

It is well documented that light has the major control of leaf economics. For example, specific leaf area, an integrative indicator of leaf structure that co-varies with leaf nitrogen content (Anten et al., 1998), shows more plastic responses to light than to nutrient availability (Poorter et al., 2009; Poorter et al., 2010). Mechanistic models can be used to interpret measured biological data (Chen et al., 2015, 2018), as in our model here,



**Fig. 5.** Effects of photosynthetic nitrogen ( $N_{ph}$ ) distributions with different values of  $f_d$  (Fig. 4) on daily canopy carbon assimilation (DCA) under different daily photosynthetic photon integrals (DPI, mol photons  $m^{-2} d^{-1}$ ) relative to average DPI during acclimation (aDPI). (A) Two-fold aDPI (aDPI<sub>200</sub>); (B) aDPI (aDPI<sub>100</sub>); (C) half aDPI (aDPI<sub>50</sub>). Acropetal  $N_{ph}$  reallocation increases with  $f_d$ . Plants grown under high nitrogen and high light (HN+HL, orange open circles), under high nitrogen and low light (HN+LL, black closed circles), under low nitrogen and high light (LN+HL, orange open triangles), and under low nitrogen and low light (LN+LL, black closed triangles) are compared under given DPI. The relative change in DCA was calculated by dividing the DCA obtained with a given  $N_{ph}$  distribution by the DCA obtained with the control  $N_{ph}$  distribution ( $f_d=1$ ) under same DPI. A change within  $\pm 5\%$  (grey shading) is considered insignificant.

providing a quantitative explanation of the different plastic responses in leaf nitrogen economics (e.g. photosynthetic nitrogen per unit leaf area,  $N_{ph}$ , and photosynthetic capacities) to light and to nitrogen by their effects on the maximum protein synthesis rate ( $S_{max}$ ; Fig. 1). A 5-fold increase in light (4–20 mol photons  $m^{-2} d^{-1}$ ) doubled  $S_{max}$  of the carboxylation pool ( $N_V$ ) and electron transport pool ( $N_J$ ; Fig. 1A, B), which is similar to the published values (Niinemets *et al.*, 2015). In contrast, increasing nitrogen supply from 2 to 10 mM increased  $S_{max}$  of  $N_V$  and  $N_J$  only by 20% and 16%, respectively. The effects of light on photosynthetic nitrogen can be quantitative (on  $N_{ph}$ ) or qualitative (on nitrogen partitioning,  $p_X$ ; Niinemets *et al.*,



**Fig. 6.** Increase in daily canopy carbon assimilation (DCA) by optimizing photosynthetic nitrogen ( $N_{ph}$ ) partitioning for different growth conditions under various daily photosynthetic photon integrals (DPI, mol photons  $m^{-2} d^{-1}$ ). The increase in DCA was the DCA with the optimal partitioning under a given DPI in comparison with the control partitioning [ $f_{p,X}=1$  in Eq. (19)]. An increase less than 5% (grey shading) is considered insignificant. The average DPI during acclimation (aDPI) is indicated by the orange arrow for HL (21.4 mol photons  $m^{-2} d^{-1}$ ) and by the black arrow for LL (8.5 mol photons  $m^{-2} d^{-1}$ ). The asterisks indicate the scenarios compared in Figs 7, 8 and Table 4 with 50%, 100% and 200% aDPI. The symbols and colors used here are the same as those in Fig. 5.

2006; Buckley *et al.*, 2013), while nitrogen only affected  $N_{ph}$  by restricting  $S_{max}$  (Figs 1, 3). Similar effects of light and nitrogen availabilities on the partitioning between electron transport and light harvesting functions were observed in spinach (Terashima and Evans, 1988). Our model of protein turnover explains the photosynthetic acclimation to light and nitrogen supply and provides a mechanistic insight into leaf nitrogen economics.


In a growing canopy, leaf age is associated with decreasing light availability (Niinemets *et al.*, 2006; Chen *et al.*, 2014). Therefore, leaf photosynthetic acclimation to light occurs together with leaf ageing, which is characterized by the protein degradation constant  $D_r$  and the constant  $t_d$  describing the decrease of protein synthesis rate in our model. The  $D_r$  values of  $N_V$  and  $N_J$  fall within the range of *in vivo* quantifications reported by Peterson *et al.* (1973) and Li *et al.* (2017). The low value of  $t_d$  (Table 1) explains the modest influence of ageing on leaf photosynthetic capacity observed under constant light conditions (Pettersen *et al.*, 2010a).

Besides light and nitrogen availability, temperature has effects on photosynthetic nitrogen content and partitioning (Yamori *et al.*, 2005; Kattge and Knorr, 2007; Yamori *et al.*, 2009). Temperature dependency of developmental processes and biochemical reactions is often described by exponential or Arrhenius-type functions (Parent *et al.*, 2010; Parent and Tardieu, 2012; Kahlen and Chen, 2015). In our model, temperature effects are considered partly by the temperature sum, which assumes a linear relationship between protein synthesis and leaf temperature. Since the exact temperature dependency of protein synthesis and degradation is unknown and our data are obtained from controlled environments with minimized temperature fluctuations, we apply the linear parsimonious approach to avoid speculation and overparameterization (Parent *et al.*, 2016).

**Table 4.** Increase in the daily canopy carbon assimilation (DCA) by optimized photosynthetic nitrogen distribution or partitioning under various daily photosynthetic photon integrals (DPI, mol photons  $\text{m}^{-2} \text{d}^{-1}$ ) for canopies grown under different conditions

Growth condition	Light level		Control DCA (mol $\text{CO}_2 \text{d}^{-1}$ )	Increase in DCA (%) by optimized	
	aDPI level (%)	DPI		Distribution	Partitioning
HN+HL	200	42.7	0.5467	<5%	6.3%
	100	21.4	0.3217	<5%	<5%
	50	10.7	0.1368	<5%	7.1%
HN+LL	200	17.1	0.2554	<5%	<5%
	100	8.5	0.1195	<5%	<5%
	50	4.3	0.0259	<5%	23.6%
LN+HL	200	42.7	0.4011	7.0%	12.7%
	100	21.4	0.2653	<5%	<5%
	50	10.7	0.1221	<5%	<5%
LN+LL	200	17.1	0.2261	6.9%	<5%
	100	8.5	0.1108	<5%	<5%
	50	4.3	0.0215	<5%	25.0%

Average DPI during acclimation (100% aDPI), 200% and 50% aDPI were tested. The increase in DCA for plants grown under the combinations of high nitrogen (HN), high light (HL), low nitrogen (LN), and low light (LL) was calculated by comparing the DCA between optimal and control distribution or partitioning.




**Fig. 7.** Ratio between optimal and control partitioning fractions (optimal  $p_x$ /control  $p_x$ ) of the carboxylation pool ( $p_v$ , orange circles), the electron transport pool ( $p_j$ , red triangles), the light harvesting pool ( $p_c$ , green squares), and contributions of daily leaf carbon assimilation (DLA) to the daily canopy carbon assimilation (DCA) increase by optimal partitioning (grey bars, right y-axis) along the canopy depth [leaf area index (LAI)  $\text{m}^2 \text{m}^{-2}$ ] under 200% average daily photosynthetic photon integral during acclimation (aDPI<sub>200</sub>) for plants grown under (A) high nitrogen and high light (HN+HL), (B) high nitrogen and low light (HN+LL), (C) low nitrogen and high light (LN+HL), (D) low nitrogen and low light (LN+LL) conditions. Photosynthetic nitrogen partitioning is close to optimum for HN+LL and LN+LL under aDPI<sub>200</sub>, which corresponds to a DPI of 42.7 and 17.1 mol photons  $\text{m}^{-2} \text{d}^{-1}$  for HL and LL, respectively. See Table 4 for the increase in DCA by the optimal partitioning.

*Above-optimum Rubisco investment can be a mechanism to adapt canopy photosynthesis to short-term light fluctuations*

Under sufficient nitrogen availability, Rubisco can function as a storage protein, which means that the amount of Rubisco can exceed the requirements to support photosynthesis

(Carmo-Silva *et al.*, 2015). The Rubisco pool has the highest value of  $k_N$  (Table 1), indicating that Rubisco synthesis reacts with higher sensitivity to increasing nitrogen availability than the other two pools. This explains the increase in the ratio between  $V_{\text{cmax}}$  and  $J_{\text{max}}$  with nitrogen availability (Hikosaka, 2004; Yamori *et al.*, 2011a), especially under LL (Fig. 3C, E).



**Fig. 8.** Ratio between optimal and control partitioning fractions (optimal  $p_{x}$ /control  $p_{x}$ ), and contributions of daily leaf carbon assimilation (DLA) to the daily canopy carbon assimilation (DCA) increase by optimal partitioning (grey bars, right y-axis) along the canopy depth [leaf area index (LAI) m<sup>2</sup> m<sup>-2</sup>] under 50% average daily photon integral during acclimation (aDPI<sub>50</sub>) for plants grown under (A) high nitrogen and high light (HN+HL), (B) high nitrogen and low light (HN+LL), (C) low nitrogen and high light (LN+HL), (D) low nitrogen and low light (LN+LL) conditions. Photosynthetic nitrogen partitioning is close to optimum for LN+HL under aDPI<sub>50</sub>, which corresponds to a DPI of 10.7 and 4.3 mol photons m<sup>-2</sup> d<sup>-1</sup> for HL and LL, respectively. The symbols and colors used here are the same as those in Fig. 7. See Table 4 for the increase in DCA by the optimal partitioning.

Under HN, Rubisco storage is advantageous since light-induced Rubisco activation, having a time constant of 3–5 min (Portis *et al.*, 1986; Kaiser *et al.*, 2018), is much faster than Rubisco synthesis. Therefore, Rubisco storage can be a mechanism for quick adaptation to a sudden increase in light. This explains why the plants grown under HN have wider ranges of DPI, at which nitrogen partitioning is optimal, than those under LN (Fig. 6; Table 4). Furthermore, excluding Rubisco activation [ $V_c = V_{cmax}$  in Eq. (9b)] in the DCA simulation resulted in a 4-fold above-optimum investment in  $N_V$  even under aDPI (data not shown). Since Rubisco is not an especially inefficient catalyst in comparison with other chemically related enzymes (Bathellier *et al.*, 2018), above-optimum Rubisco investment in the canopy can be rather a mechanism for adapting to short-term light fluctuation than a mechanism to overcome its enzymatic inefficiency.

#### Implications for crop model improvement and greenhouse management

Using plant models to understand crop performance requires knowledge of physiological mechanisms (Boote *et al.*, 2013; Poorter *et al.*, 2013). By integrating the known biological mechanism of protein turnover at the leaf level into a multi-layer model of canopy photosynthesis, we demonstrate the explanatory power of a mechanistic model for the measured biological data. Our simulations suggest that canopy photosynthesis can be increased by manipulating the functional pools

related to photosynthesis. For example, investment in Rubisco and electron transport (Ishimaru *et al.*, 2001; Yamori *et al.*, 2011b) should be increased under increasing light (Fig. 7), and a larger antenna size for light harvesting (Masuda *et al.*, 2003) is required under decreasing light availability (Fig. 8). It is clear that the pattern of optimal nitrogen partitioning depends strongly on light regime, and biosynthetic regulation is unlikely to keep up with daily light fluctuation (up to 4-fold difference; Supplementary Fig. S3).

In greenhouse cultivation, it is possible to achieve a more stable light environment using supplemental lighting. This can be a plausible solution to improve the vertical light distribution (Lu and Mitchell, 2016) and to minimize the suboptimality in nitrogen use induced by light fluctuation. Since carbon assimilation is the rate-limiting step for yield production of cucumber plants due to the indeterminate production of vegetative and generative organs (Wiechers *et al.*, 2011), canopy carbon gain can be considered as an approximation for yield. Our simulation suggests that the suboptimal nitrogen partitioning induced by a 50% decrease in DPI can be compensated by reducing the light limitation of the shaded leaves using inter-row lighting during the high-light season (ca. 7% increase in DCA) and using top-lighting, possibly in combination with inter-lighting, during the low-light season (ca. 25% increase in DCA; Table 4; Fig. 8), similar to the reported increase in cucumber fruit yield (22%–31%) by inter-lighting in the winter season (Kumar *et al.*, 2016). In the summer season, suboptimal nitrogen partitioning

induced by sudden doubling in DPI can be overcome by pre-treatment of increasing nitrogen supply and inter-lighting (ca. 6% increase in DCA; Table 4), which maintains the biochemical capacity and reduces the biochemical limitation of the shaded leaves (Pettersen *et al.*, 2010b; Trouwborst *et al.*, 2010; Chen *et al.*, 2014). These results provide a physiological explanation at canopy level for the observations of supplemental lighting experiments (Hovi *et al.*, 2004; Hovi-Pekkanen and Tahvonen, 2008; Pettersen *et al.*, 2010b; Trouwborst *et al.*, 2010). Furthermore, the relationship between protein synthesis rate and intercepted light intensity is non-linear in our model [Eq. (7)], which may offer an explanation why the photoacclimatory responses of a leaf grown under natural within-day light fluctuation differ from that under constant light, as shown in a recent experimental study (Violet-Chabrand *et al.*, 2017).

Light fluctuations occur particularly in the lower canopy layer, where sunflecks cause strong and frequent variations in light, thereby increasing variations of  $N_{ph}$  and  $p_X$  in the older leaves (Fig. 3). Interestingly, leaves under HL seemed to prioritize their nitrogen investment in  $N_j$  over  $N_C$  under LN with increasing leaf age (Fig. 3E), which might be explained by the reduced LAI development under LN+HL and, hence, the higher light interception of the older leaves (Figs S4C, S5). Since within-leaf and within-day light heterogeneity (e.g. sunflecks) were not described in the model, these variations observed in the greenhouse experiment could not be reproduced in the simulations (Supplementary Fig. S6). This can be improved by coupling the model with a 3D structural plant model and the use of shorter time steps in the simulations to capture more realistic response of photoacclimation.

## Conclusions

We propose a mechanistic model to quantify the effects of leaf age, nitrogen and light availabilities on photosynthetic acclimation. The model predicts the observed photosynthetic acclimation under different combinations of nitrogen supply and light availability in the greenhouse. Model simulation indicates that photosynthetic nitrogen distribution is close to optimum and photosynthetic nitrogen partitioning can be optimal under constant light conditions. However, large fluctuation in light between days under natural conditions inevitably leads to sub-optimal nitrogen partitioning. Our study provides insights into photosynthetic acclimation and the model can be used for crop model improvement and provides guidelines for greenhouse management.

## Supplementary data

Supplementary data are available at *JXB* online.

Fig. S1. Schematic diagram of photosynthetic nitrogen turnover.

Fig. S2. Relationship between relative chlorophyll content and leaf chlorophyll concentration.

Fig. S3. Environmental input for the model evaluation and simulation.

Fig. S4. Relationships between leaf angle, LAI, and age.

Fig. S5. Leaf area distribution used as input in the daily canopy assimilation simulation.

Fig. S6. Comparisons of simulated photosynthetic nitrogen traits between nitrogen supply levels and between light conditions.

Fig. S7. Leaf photosynthetic nitrogen distributions with the vertical light distribution.

Table S1. Canopy characteristics used in the daily canopy assimilation simulation.

## Acknowledgements

This work was supported by Deutsche Forschungsgemeinschaft (DFG). We thank Ilona Napp, Marlies Lehmann, Adjoa Sekyi-Appiah, Sanzida Akhter Anee and Felliessia Regina Halim for their assistance during the experiments.

## References

- Anten NP, Miyazawa K, Hikosaka K, Nagashima H, Hirose T. 1998. Leaf nitrogen distribution in relation to leaf age and photon flux density in dominant and subordinate plants in dense stands of a dicotyledonous herb. *Oecologia* **113**, 314–324.
- Anten NP, Schieving F, Werger MJ. 1995. Patterns of light and nitrogen distribution in relation to whole canopy carbon gain in  $C_3$  and  $C_4$  mono- and dicotyledonous species. *Oecologia* **101**, 504–513.
- Athanasίου K, Dyson BC, Webster RE, Johnson GN. 2010. Dynamic acclimation of photosynthesis increases plant fitness in changing environments. *Plant Physiology* **152**, 366–373.
- Bathellier C, Tcherkez G, Lorimer GH, Farquhar GD. 2018. Rubisco is not really so bad. *Plant, Cell & Environment* **41**, 705–716.
- Boote KJ, Jones JW, White JW, Asseng S, Lizaso JI. 2013. Putting mechanisms into crop production models. *Plant, Cell & Environment* **36**, 1658–1672.
- Buckley TN, Cescatti A, Farquhar GD. 2013. What does optimization theory actually predict about crown profiles of photosynthetic capacity when models incorporate greater realism? *Plant, Cell & Environment* **36**, 1547–1563.
- Carmo-Silva E, Scales JC, Madgwick PJ, Parry MA. 2015. Optimizing Rubisco and its regulation for greater resource use efficiency. *Plant, Cell & Environment* **38**, 1817–1832.
- Chen TW, Henke M, de Visser PH, Buck-Sorlin G, Wiechers D, Kahlen K, Stützel H. 2014. What is the most prominent factor limiting photosynthesis in different layers of a greenhouse cucumber canopy? *Annals of Botany* **114**, 677–688.
- Chen TW, Nguyen TM, Kahlen K, Stützel H. 2015. High temperature and vapor pressure deficit aggravate architectural effects but ameliorate non-architectural effects of salinity on dry mass production of tomato. *Frontiers in Plant Science* **6**, 887.
- Chen TW, Stützel H, Kahlen K. 2018. High light aggravates functional limitations of cucumber canopy photosynthesis under salinity. *Annals of Botany* **121**, 797–807.
- De Kauwe MG, Lin YS, Wright IJ, *et al.* 2016. A test of the 'one-point method' for estimating maximum carboxylation capacity from field-measured, light-saturated photosynthesis. *New Phytologist* **210**, 1130–1144.
- Dreccer MF, van Oijen M, Schapendonk AHCM, Pot CS, Rabbinge R. 2000. Dynamics of vertical leaf nitrogen distribution in a vegetative wheat canopy. Impact on canopy photosynthesis. *Annals of Botany* **86**, 821–831.
- Evans JR. 1989. Partitioning of nitrogen between and within leaves grown under different irradiances. *Functional Plant Biology* **16**, 533–548.
- Evans JR. 1993. Photosynthetic acclimation and nitrogen partitioning within a lucerne canopy. II. Stability through time and comparison with a theoretical optimum. *Functional Plant Biology* **20**, 69–82.
- Falster DS, Westoby M. 2003. Leaf size and angle vary widely across species: what consequences for light interception? *New Phytologist* **158**, 509–525.

- Farquhar GD, von Caemmerer S, Berry JA.** 1980. A biochemical model of photosynthetic CO<sub>2</sub> assimilation in leaves of C<sub>3</sub> species. *Planta* **149**, 78–90.
- Field C.** 1983. Allocating leaf nitrogen for the maximization of carbon gain: Leaf age as a control on the allocation program. *Oecologia* **56**, 341–347.
- Flexas J, Ribas-Carbó M, Diaz-Espejo A, Galmés J, Medrano H.** 2008. Mesophyll conductance to CO<sub>2</sub>: current knowledge and future prospects. *Plant, Cell & Environment* **31**, 602–621.
- Harley PC, Loreto F, Di Marco G, Sharkey TD.** 1992. Theoretical considerations when estimating the mesophyll conductance to CO<sub>2</sub> flux by analysis of the response of photosynthesis to CO<sub>2</sub>. *Plant Physiology* **98**, 1429–1436.
- Hikosaka K.** 2004. Interspecific difference in the photosynthesis-nitrogen relationship: patterns, physiological causes, and ecological importance. *Journal of Plant Research* **117**, 481–494.
- Hikosaka K.** 2005. Nitrogen partitioning in the photosynthetic apparatus of *Plantago asiatica* leaves grown under different temperature and light conditions: similarities and differences between temperature and light acclimation. *Plant & Cell Physiology* **46**, 1283–1290.
- Hikosaka K.** 2014. Optimal nitrogen distribution within a leaf canopy under direct and diffuse light. *Plant, Cell & Environment* **37**, 2077–2085.
- Hikosaka K.** 2016. Optimality of nitrogen distribution among leaves in plant canopies. *Journal of Plant Research* **129**, 299–311.
- Hikosaka K, Anten NP, Borjigidai A, et al.** 2016. A meta-analysis of leaf nitrogen distribution within plant canopies. *Annals of Botany* **118**, 239–247.
- Hikosaka K, Terashima I.** 1996. Nitrogen partitioning among photosynthetic components and its consequence in sun and shade plants. *Functional Ecology* 335–343.
- Hirose T, Ackerly DD, Traw MB, Ramseier D, Bazzaz FA.** 1997. CO<sub>2</sub> elevation, canopy photosynthesis, and optimal leaf area index. *Ecology* **78**, 2339–2350.
- Hirose T, Werger MJ.** 1987. Maximizing daily canopy photosynthesis with respect to the leaf nitrogen allocation pattern in the canopy. *Oecologia* **72**, 520–526.
- Hollinger DY.** 1996. Optimality and nitrogen allocation in a tree canopy. *Tree Physiology* **16**, 627–634.
- Hovi T, Näskilä J, Tahvonen R.** 2004. Interlighting improves production of year-round cucumber. *Scientia Horticulturae* **102**, 283–294.
- Hovi-Pekkanen T, Tahvonen R.** 2008. Effects of interlighting on yield and external fruit quality in year-round cultivated cucumber. *Scientia Horticulturae* **116**, 152–161.
- Irving LJ, Robinson D.** 2006. A dynamic model of Rubisco turnover in cereal leaves. *New Phytologist* **169**, 493–504.
- Ishimaru K, Kobayashi N, Ono K, Yano M, Ohsugi R.** 2001. Are contents of Rubisco, soluble protein and nitrogen in flag leaves of rice controlled by the same genetics? *Journal of Experimental Botany* **52**, 1827–1833.
- Kahlen K, Chen TW.** 2015. Predicting plant performance under simultaneously changing environmental conditions—The interplay between temperature, light, and internode growth. *Frontiers in Plant Science* **6**, 1130.
- Kahlen K, Stützel H.** 2011. Modelling photo-modulated internode elongation in growing glasshouse cucumber canopies. *New Phytologist* **190**, 697–708.
- Kaiser E, Morales A, Harbinson J.** 2018. Fluctuating light takes crop photosynthesis on a rollercoaster ride. *Plant Physiology* **176**, 977–989.
- Kattge J, Knorr W.** 2007. Temperature acclimation in a biochemical model of photosynthesis: a reanalysis of data from 36 species. *Plant, Cell & Environment* **30**, 1176–1190.
- Kimball BA, Bellamy LA.** 1986. Generation of diurnal solar radiation, temperature, and humidity patterns. *Energy in Agriculture* **5**, 185–197.
- Kitao M, Kitaoka S, Harayama H, Tobita H, Agathokleous E, Utsugi H.** 2018. Canopy nitrogen distribution is optimized to prevent photoinhibition throughout the canopy during sun flecks. *Scientific Reports* **8**, 503.
- Kok B.** 1948. A critical consideration of the quantum yield of *Chlorella* photosynthesis. *Enzymologia* **13**, 1–56.
- Kumar K, Hao X, Khosla S, Guo X, Bennett N.** 2016. Comparison of HPS lighting and hybrid lighting with top HPS and intra-canopy LED lighting for high-wire mini-cucumber production. *Acta Horticulturae* **1134**, 111–118.
- Li L, Nelson CJ, Trösch J, Castleden I, Huang S, Millar AH.** 2017. Protein degradation rate in *Arabidopsis thaliana* leaf growth and development. *The Plant Cell* **29**, 207–228.
- Lichtenthaler HK.** 1987. Chlorophylls and carotenoids: pigments of photosynthetic biomembranes. *Methods in Enzymology* **148**, 350–383.
- Loriaux SD, Avenson TJ, Welles JM, McDermitt DK, Eckles RD, Riensche B, Genty B.** 2013. Closing in on maximum yield of chlorophyll fluorescence using a single multiphase flash of sub-saturating intensity. *Plant, Cell & Environment* **36**, 1755–1770.
- Lu N, Mitchell CA.** 2016. Supplemental lighting for greenhouse-grown fruiting vegetables. In: Kozai T, Fujiwara K, Runkle ES, eds. *LED lighting for urban agriculture*. Singapore: Springer, 219–232.
- Masuda T, Tanaka A, Melis A.** 2003. Chlorophyll antenna size adjustments by irradiance in *Dunaliella salina* involve coordinate regulation of chlorophyll a oxygenase (CAO) and Lhcb gene expression. *Plant Molecular Biology* **51**, 757–771.
- Medlyn BE, Duursma RA, Eamus D, Ellsworth DS, Prentice IC, Barton CVM, Crous KY, De Angelis P, Freeman M, Wingate L.** 2011. Reconciling the optimal and empirical approaches to modelling stomatal conductance. *Global Change Biology* **17**, 2134–2144.
- Meir P, Kruijt B, Broadmeadow M, Barbosa E, Kull O, Carswell F, Nobre A, Jarvis PG.** 2002. Acclimation of photosynthetic capacity to irradiance in tree canopies in relation to leaf nitrogen concentration and leaf mass per unit area. *Plant, Cell & Environment* **25**, 343–357.
- Monsi M, Saeki T.** 1953. Über den Lichtfaktor in den Pflanzengesellschaften und seine Bedeutung für die Stoffproduktion. *Japanese Journal of Botany* **14**, 22–52. [Republished in English: Monsi M, Saeki T. 2005. On the factor light in plant communities and its importance for matter production. *Annals of Botany* 95, 549–567.]
- Moreau D, Allard V, Gaju O, Le Gouis J, Foulkes MJ, Martre P.** 2012. Acclimation of leaf nitrogen to vertical light gradient at anthesis in wheat is a whole-plant process that scales with the size of the canopy. *Plant Physiology* **160**, 1479–1490.
- Moualeu-Ngangue DP, Chen TW, Stützel H.** 2016. A modeling approach to quantify the effects of stomatal behavior and mesophyll conductance on leaf water use efficiency. *Frontiers in Plant Science* **7**, 875.
- Moualeu-Ngangue DP, Chen TW, Stützel H.** 2017. A new method to estimate photosynthetic parameters through net assimilation rate–intercellular space CO<sub>2</sub> concentration (A–C<sub>i</sub>) curve and chlorophyll fluorescence measurements. *New Phytologist* **213**, 1543–1554.
- Nelson DW, Sommers LE.** 1980. Total nitrogen analysis of soil and plant tissues. *Journal of the Association of Official Analytical Chemists* **63**, 770–778.
- Niinemets Ü.** 2016. Leaf age dependent changes in within-canopy variation in leaf functional traits: a meta-analysis. *Journal of Plant Research* **129**, 313–338.
- Niinemets U, Cescatti A, Rodeghiero M, Tosens T.** 2006. Complex adjustments of photosynthetic potentials and internal diffusion conductance to current and previous light availabilities and leaf age in Mediterranean evergreen species *Quercus ilex*. *Plant, Cell & Environment* **29**, 1159–1178.
- Niinemets Ü, Keenan TF, Hallik L.** 2015. A worldwide analysis of within-canopy variations in leaf structural, chemical and physiological traits across plant functional types. *New Phytologist* **205**, 973–993.
- Ögren E, Evans JR.** 1993. Photosynthetic light-response curves. *Planta* **189**, 182–190.
- Parent B, Tardieu F.** 2012. Temperature responses of developmental processes have not been affected by breeding in different ecological areas for 17 crop species. *New Phytologist* **194**, 760–774.
- Parent B, Turc O, Gibon Y, Stitt M, Tardieu F.** 2010. Modelling temperature-compensated physiological rates, based on the co-ordination of responses to temperature of developmental processes. *Journal of Experimental Botany* **61**, 2057–2069.
- Parent B, Vile D, Violle C, Tardieu F.** 2016. Towards parsimonious ecophysiological models that bridge ecology and agronomy. *New Phytologist* **210**, 380–382.
- Peterson LW, Kleinkopf GE, Huffaker RC.** 1973. Evidence for lack of turnover of ribulose 1,5-diphosphate carboxylase in barley leaves. *Plant Physiology* **51**, 1042–1045.

- Pettersen RI, Torre S, Gisleørud HR.** 2010a. Effects of leaf aging and light duration on photosynthetic characteristics in a cucumber canopy. *Scientia Horticulturae* **125**, 82–87.
- Pettersen RI, Torre S, Gisleørud HR.** 2010b. Effects of intracanal lighting on photosynthetic characteristics in cucumber. *Scientia Horticulturae* **125**, 77–81.
- Pons TL, Anten NP.** 2004. Is plasticity in partitioning of photosynthetic resources between and within leaves important for whole-plant carbon gain in canopies? *Functional Ecology* **18**, 802–811.
- Poorter H, Anten NP, Marcelis LF.** 2013. Physiological mechanisms in plant growth models: do we need a supra-cellular systems biology approach? *Plant, Cell & Environment* **36**, 1673–1690.
- Poorter H, Niinemets U, Poorter L, Wright IJ, Villar R.** 2009. Causes and consequences of variation in leaf mass per area (LMA): a meta-analysis. *New Phytologist* **182**, 565–588.
- Poorter H, Niinemets U, Walter A, Fiorani F, Schurr U.** 2010. A method to construct dose–response curves for a wide range of environmental factors and plant traits by means of a meta-analysis of phenotypic data. *Journal of Experimental Botany* **61**, 2043–2055.
- Portis AR, Salvucci ME, Ogren WL.** 1986. Activation of ribulosebiphosphate carboxylase/oxygenase at physiological CO<sub>2</sub> and ribulosebiphosphate concentrations by rubisco activase. *Plant Physiology* **82**, 967–971.
- Prieto JA, Louarn G, Perez Peña J, Ojeda H, Simonneau T, Lebon E.** 2012. A leaf gas exchange model that accounts for intra-canopy variability by considering leaf nitrogen content and local acclimation to radiation in grapevine (*Vitis vinifera* L.). *Plant, Cell & Environment* **35**, 1313–1328.
- Qian T, Elings A, Dieleman JA, Gort G, Marcelis LF.** 2012. Estimation of photosynthesis parameters for a modified Farquhar-von Caemmerer-Berry model using simultaneous estimation method and nonlinear mixed effects model. *Environmental and Experimental Botany* **82**, 66–73.
- Retkute R, Smith-Unna SE, Smith RW, Burgess AJ, Jensen OE, Johnson GN, Preston SP, Murchie EH.** 2015. Exploiting heterogeneous environments: does photosynthetic acclimation optimize carbon gain in fluctuating light? *Journal of Experimental Botany* **66**, 2437–2447.
- Savvides A, Dieleman JA, van Ieperen W, Marcelis LF.** 2016. A unique approach to demonstrating that apical bud temperature specifically determines leaf initiation rate in the dicot *Cucumis sativus*. *Planta* **243**, 1071–1079.
- Singsaas EL, Ort DR, Delucia EH.** 2004. Elevated CO<sub>2</sub> effects on mesophyll conductance and its consequences for interpreting photosynthetic physiology. *Plant, Cell & Environment* **27**, 41–50.
- Song Q, Wang Y, Qu M, Ort DR, Zhu XG.** 2017. The impact of modifying photosystem antenna size on canopy photosynthetic efficiency—Development of a new canopy photosynthesis model scaling from metabolism to canopy level processes. *Plant, Cell & Environment* **40**, 2946–2957.
- Suzuki Y, Makino A, Mae T.** 2001. Changes in the turnover of Rubisco and levels of mRNAs of *rbcL* and *rbcS* in rice leaves from emergence to senescence. *Plant, Cell & Environment* **24**, 1353–1360.
- Terashima I, Evans JR.** 1988. Effects of light and nitrogen nutrition on the organization of the photosynthetic apparatus in spinach. *Plant and Cell Physiology* **29**, 143–155.
- Thornley JHM.** 1998. Dynamic model of leaf photosynthesis with acclimation to light and nitrogen. *Annals of Botany* **81**, 421–430.
- Trouwborst G, Hogewoning SW, Harbinson J, van Ieperen W.** 2011. Photosynthetic acclimation in relation to nitrogen allocation in cucumber leaves in response to changes in irradiance. *Physiologia Plantarum* **142**, 157–169.
- Trouwborst G, Oosterkamp J, Hogewoning SW, Harbinson J, van Ieperen W.** 2010. The responses of light interception, photosynthesis and fruit yield of cucumber to LED-lighting within the canopy. *Physiologia Plantarum* **138**, 289–300.
- Verkroost AW, Wassen MJ.** 2005. A simple model for nitrogen-limited plant growth and nitrogen allocation. *Annals of Botany* **96**, 871–876.
- Violet-Chabrand S, Matthews JS, Simkin AJ, Raines CA, Lawson T.** 2017. Importance of fluctuations in light on plant photosynthetic acclimation. *Plant Physiology* **173**, 2163–2179.
- Walters RG.** 2005. Towards an understanding of photosynthetic acclimation. *Journal of Experimental Botany* **56**, 435–447.
- Werger MJ, Hirose T.** 1991. Leaf nitrogen distribution and whole canopy photosynthetic carbon gain in herbaceous stands. *Vegetatio* **97**, 11–20.
- Wiechers D, Kahlen K, Stützel H.** 2011. Dry matter partitioning models for the simulation of individual fruit growth in greenhouse cucumber canopies. *Annals of Botany* **108**, 1075–1084.
- Wilson KB, Baldocchi DD, Hanson PJ.** 2000. Spatial and seasonal variability of photosynthetic parameters and their relationship to leaf nitrogen in a deciduous forest. *Tree Physiology* **20**, 565–578.
- Wright IJ, Leishman MR, Read C, Westoby M.** 2006. Gradients of light availability and leaf traits with leaf age and canopy position in 28 Australian shrubs and trees. *Functional Plant Biology* **33**, 407–419.
- Yamori W, Evans JR, Von Caemmerer S.** 2010. Effects of growth and measurement light intensities on temperature dependence of CO<sub>2</sub> assimilation rate in tobacco leaves. *Plant, Cell & Environment* **33**, 332–343.
- Yamori W, Nagai T, Makino A.** 2011a. The rate-limiting step for CO<sub>2</sub> assimilation at different temperatures is influenced by the leaf nitrogen content in several C<sub>3</sub> crop species. *Plant, Cell & Environment* **34**, 764–777.
- Yamori W, Noguchi KO, Terashima I.** 2005. Temperature acclimation of photosynthesis in spinach leaves: analyses of photosynthetic components and temperature dependencies of photosynthetic partial reactions. *Plant, Cell & Environment* **28**, 536–547.
- Yamori W, Noguchi K, Hikosaka K, Terashima I.** 2009. Cold-tolerant crop species have greater temperature homeostasis of leaf respiration and photosynthesis than cold-sensitive species. *Plant & Cell Physiology* **50**, 203–215.
- Yamori W, Takahashi S, Makino A, Price GD, Badger MR, von Caemmerer S.** 2011b. The roles of ATP synthase and the cytochrome b6/f complexes in limiting chloroplast electron transport and determining photosynthetic capacity. *Plant Physiology* **155**, 956–962.
- Zhu XG, Long SP, Ort DR.** 2010. Improving photosynthetic efficiency for greater yield. *Annual Review of Plant Biology* **61**, 235–261.